



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

A61K 49/00

A2

(11) International Publication Number:

WO 92/17214

(43) International Publication Date:

15 October 1992 (15.10.92)

(21) International Application Number: PCT/US92/02685

(22) International Filing Date: 3 April 1992 (03.04.92)

(30) Priority data:

680,675

4 April 1991 (04.04.91)


US

(71)(72) Applicant and Inventor: RANNEY, David, F. [US/US];  
3539 Courtdale Drive, Dallas, TX 75234 (US).(74) Agent: HODGINS, Daniel, S.; Arnold, White & Durkee,  
P.O. Box 4433, Houston, TX 77210 (US).(81) Designated States: AT, AT (European patent), AU, BB, BE  
(European patent), BF (OAPI patent), BG, BJ (OAPI pa-  
tent), BR, CA, CF (OAPI patent), CG (OAPI patent),  
CH, CH (European patent), CI (OAPI patent), CM (OA-  
PI patent), CS, DE, DE (European patent), DK, DK (Eu-  
ropean patent), ES, ES (European patent), FI, FR (Euro-  
pean patent), GA (OAPI patent), GB, GB (European pa-  
tent), GN (OAPI patent), GR (European patent), HU, IT  
(European patent), JP, KP, KR, LK, LU, LU (European  
patent), MC (European patent), MG, ML (OAPI patent),  
MN, MR (OAPI patent), MW, NL, NL (European pa-  
tent), NO, PL, RO, RU, SD, SE, SE (European patent),  
SN (OAPI patent), TD (OAPI patent), TG (OAPI pa-  
tent).

Published

*Without international search report and to be republished  
upon receipt of that report.*

(54) Title: PHYSICALLY AND CHEMICALLY STABILIZED POLYATOMIC CLUSTERS FOR MAGNETIC RESONANCE IMAGE AND SPECTRAL ENHANCEMENT



## (57) Abstract

Improved compositions and methods for selective access to tumor regions (or other regions of abnormal endothelial properties). This capability provides powerful contrast-enhancement agents for nuclear magnetic resonance imaging. A polyatomic complex which includes intramolecular ferromagnetic coupling between metal atoms is associated with a polymer or microsphere carrier matrix which will bind to endothelial determinants. A solution containing this carrier complex is injected into a human (or other) body to be imaged. The carrier complex will preferentially extravasate at locations where the blood vessel walls have increased porosity or microvascular surface changes, and especially at tumor sites. Thus, the changes in relaxation time induced by the presence of the carrier complex will provide a high-gain marker for magnetic resonance imaging. Multiple superparamagnetic polyatomic complexes are described, including novel complexes which include acetate and glycinate bridging ligands with a polyatomic metal-atom-complex core.

BEST AVAILABLE COPY

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Tchad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

**PHYSICALLY AND CHEMICALLY STABILIZED POLYATOMIC CLUSTERS  
FOR MAGNETIC RESONANCE IMAGE AND SPECTRAL ENHANCEMENT**

5

The present application relates to formulations of polyatomic clusters which are rendered useful for in vivo magnetic resonance imaging (MRI) and spectral enhancement by means of physical, chemical, and combined physical-chemical stabilization. Said clusters are defined as polyatomic molecular complexes containing multiple atoms of either the same or of different atomic number, wherein two or more of the atoms have a magnetic moment. In selected cases, the polyatomic cluster may cause the atoms of magnetic moment to be oriented in ferromagnetic alignment, such that their multiple magnetic moments are additive intramolecularly, thereby giving the cluster a total magnetic moment greater than that of any individual atom of magnetic moment. Said polyatomic clusters may contain single or plural bridging ligands (for the purposes of this disclosure the term "bridging ligand" is equivalent to the term "bridging molecular or atomic species"), however, such bridging ligands may be distinguished from classical chelators, in that the bridging ligands, when plural, are not covalently linked together to form multiple covalent coordination sites.

Stabilization of said polyatomic clusters by the present compositions and methods of formulation results in favorable potency, plasma stability, biodistribution, delivery to desired target sites, excretion, toxicity, and/or other favorable in vivo properties, in comparison to parallel properties of the native polyatomic clusters alone. Said polyatomic cluster formulations include, but are not limited to clusters which are chemically or physically associated with carrier substances. Carrier substances are defined as substances with which the

polyatomic clusters are chemically or physically associated for the purpose of altering the in vivo properties described above. In this context, such carrier substances are distinguished from existing, traditional formulation carriers and pharmaceutical vehicles, in that these existing, traditional substances are used principally to render diagnostic agents in optimal states for solubilization, processing, drying or other aspects of in vitro formulation.

10

Association and stabilization of the present polyatomic clusters with carrier substances may be by physical, chemical or combined physical-chemical means. Chemical means may include, but are not limited to binding of said clusters to carrier substances by hydrophobic bonding, hydrogen bonding, paired-ion association, multiple paired-ion association, and partial or total coordination or trans-chelation binding of the polyatomic clusters to the carrier substances via single or multiple ligands which are themselves bound to these carrier substances. In the case of coordination or trans-chelation binding of polyatomic clusters to carrier substances, it is important to note that such coordination or trans-chelation is explicitly for the purpose of intermolecular attachment of the polyatomic cluster to the carrier substance, and not for the purpose of intramolecular stabilization of the polyatomic cluster. Hence, although partial chelation of the polyatomic cluster may result as a byproduct of carrier binding, the polyatomic cluster is considered to remain as such and not to change into a classical (full) chelate by virtue of this mechanism of carrier binding.

The formulations used for physical and combined physical-chemical stabilization typically include one or more of: emulsification, dispersion stabilization, low or

35



high pressure homogenization, sonification, heat stabilization, other energy-input processes, organic solvent processing, aqueous solvent processing, combined organic-aqueous cosolvent processing, cosolvent processing wherein one or more of the cosolvents is a nonsolvent or partial nonsolvent for the polyatomic cluster or the carrier substance or both, and other forms of chemical stabilization or binding.

10       Magnetic resonance images (MRI) have been obtained from the brain and body of living animals and humans by means of applying pulsed radio frequency (rf) energy in the presence of strong fixed and gradient magnetic fields, and monitoring the induced signals given off by  
15       atoms and ions with a magnetic moment and characteristic frequency of nuclear oscillation in the defined strong magnetic field. One major advantage of MRI images, in relation to other medical imaging modalities, is that they provide an important means of clinical brain and  
20       body imaging of internal structures, organs and foci of disease typically at a very high spatial resolution of less than about 0.3 x 0.3 mm in-plane resolution and 2 mm slice thickness.

25       MRI image enhancement has been accomplished successfully in vivo by producing alterations in the relaxation times and/or spectral shifts of paramagnetic atoms present in the animal or human being imaged. Such enhancements (spectral shifts) have been produced by  
30       administering, among others, the following major classes and examples of MRI diagnostic agents:

1.       paramagnetic ions and chelates, such as gadolinium diethylenetriamine pentaacetic acid dimeglumine (Gd-DTPA dimeglumine -- Berlex-Schering AG; West Germany patent filed by  
35

Gries, Rosenberg and Weinman: DE-OS 3129906 A 1 (1981)), and gadolinium DOTA;

2. superparamagnetic substances, such as dextran iron;
3. ferromagnetic substances, such as micronized iron; and
4. polyatomic molecular clusters, including:
  - a. (1) homopolyatomic clusters, including the  $\text{Cr}^{\text{III}}$  cluster,  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  (A Bino, et al, US Patent No. 4,832,877, issued May 23, 1989; and A Bino, et al, Science, September 16, 1988); and
  - a. (2) heteropolyatomic clusters.

In clinical imaging, these substances typically produce enhancement by altering the T1 or T2 relaxation times (or shifting the spectra) of induced magnetic signals given off by paramagnetic atoms present in the body, usually comprising water and fat protons and less commonly including carbon and phosphorous nuclei, but potentially including exogenously administered magnetic ions and atoms such as  $^{19}\text{fluorine}$ . Paramagnetics typically act as T1-weighted enhancing agents. Superparamagnetics and ferromagnetics typically act as T2-weighted enhancing agents. Polyatomic clusters may act as either T1- or T2-weighted agents depending on their size; the orientation of their intramolecular component atomic magnetic moments; and the activity or cancellation of intermolecular magnetic moments, which may be influenced by various super-molecular formulation states.

-5-

The major difficulties encountered in discovering and formulating previous MRI image enhancing (spectral shift) agents are as follows:

- 5           1.   these agents are typically quite toxic in their  
            free ionic form -- particularly for the  
            currently favored, paramagnetic ion,  
            gadolinium, which is the most potent ion  
10           available for paramagnetic relaxation due to  
            its maximal number of seven unpaired electrons;
2.   the free agents are typically water insoluble  
                or very poorly water soluble at physiologic pH;
- 15           3.   the agents must be administered in relatively  
                high doses due to the inherent insensitivity of  
                MRI imaging;
4.   after chelation of the existing paramagnetic  
20           agents, which is required to confer acceptable  
                water solubility, toxicity and body clearance,  
                these agents:
  - 25           a.   lose a significant fraction of their  
                potency because the chelator's functional  
                groups occupy almost all of the  
                paramagnetic ion's inner coordination  
                sites coinciding with the strongest  
                portion of the ion's paramagnetic dipole;  
                and
  - 30           b.   act as nonselective (nonspecific) contrast  
                agents due to their rapid free diffusion  
                into normal extracellular body fluid, as  
                well as into sites of disease (Note: in  
35           the brain but not the body, such diffusion  
                into the normal extracellular fluid is

-6-

reduced but not entirely eliminated by the presence of a blood-brain barrier);

5. due to their rapid free diffusion in body water, previous paramagnetic-ion chelates:
  - a. typically fail to enhance the very small lesions (sites of disease less than 5 mm in diameter), which are theoretically capable of resolution by current MRI instrumentation technology;
  - b. fail to optimally discriminate between perfused and nonperfused subregions of tumor nodules and localized infections, due to the their even more rapid diffusion through the less structured (bound) interstitial water (extracellular tissue compartment) within these sites of disease; and
  - c. backdiffuse very rapidly from these sites of disease into the bloodstream, thereby minimizing the time available for image acquisition after intravenous administration of these contrast agents (typically to less than 15-30 minutes), thereby eliminating the possibility of pre-dosing patients outside the imaging room.
6. existing superparamagnetic and ferromagnetic agents:
  - a. require a relatively long time for total body clearance (typically several days to weeks), thereby presenting substantial regulatory disadvantages; and
  - b. act predominantly as T2-weighted agents, which typically darken rather than

-7-

brighten (enhance) MRI images of the tissues and sites in which these agents become localized.

- 5           7. existing paramagnetic, superparamagnetic and ferromagnetic agents are either ineffective or markedly suboptimal for MRI blood-pool and perfusion image contrast enhancement.

10           It has previously been shown by the present applicant, as disclosed in previous filings (International Applications PCT/US88/01096, PCT/US89/04295, that selectivity of uptake at sites of disease could be achieved for MRI contrast agents by  
15 simultaneously employing both of the mechanisms described in Nos. 1. and 2. (immediately below):

- 20           1. Covalent or noncovalent association of magnetic agents with carrier substances which confer sufficient size -- at least about 15,000 to 20,000 MW -- to exclude rapid, free exchange of diagnostic agents between the blood compartment and the extracellular fluid compartment (interstitium) of normal body tissues; and  
25
- 30           2. Covalent and noncovalent association of magnetic agents with carrier substances which bind selectively to determinants induced on the endothelium (and epithelium) at sites of disease; and normally present on the endothelium of certain organs (e.g., lung among others). Such binding, in turn, induces active uptake of the agent-carrier combination across these intermediate barrier structures  
35 (endothelium and epithelium) into the interstitial compartment of the diseased

tissues (lesional sites) and in certain cases, normal target organs. For particular carrier substances, including but not limited to heparins and heparan sulfates, such selective, active lesional uptake has also been shown to be accompanied by prolonged lesional retention of the diagnostic-carrier (or drug-carrier) combination. This is due theoretically to binding of heparins (and other related, usually negatively charged carbohydrates and glycoaminoglycans) to endogenous heparan sulfates and related binding substituents which become induced or exposed within the interstitial compartment of tumors, infections, tissues altered by various inflammatory conditions and other lesional sites, particularly those involving vascular and/or tissue remodeling as a component of the disease.

Among the novel teachings set forth in the present application is the recent unpublished data obtained by the present applicant, which indicate that the active uptake of heparin, and potentially other negatively charged carbohydrates and related molecules, occurs selectively at lesional sites even below the applicants' and others' previously published size-exclusion cutoff of 15,000 to 20,000 Daltons, and hence, occurs even in the absence of the size exclusion mechanism described in No. 1, immediately above. This affords a significantly improved basis for site-selective uptake of carrier-associated diagnostic (and therapeutic) agents in tumors and other lesional sites. It also provides a new basis for constructing novel, selective MRI contrast agents, including polyatomic cluster-carrier compositions, at a size substantially less than the approximately 15,000 MW

-9-

cutoff at which size exclusion prevents their rapid, free exchange into normal body tissues. Such novel MRI contrast agents have multiple advantages, including:

- 5           1.   the most rapid plasma clearance available for  
any site- selective agent described to date ( $t_{1/2}$  of less than about 30 minutes, which very  
nearly equals the  $t_{1/2}$ 's of 20 minutes  
characteristic of nonselective agents),  
10           producing reduced background enhancement from  
plasma and para-lesional sites at early  
postinjection imaging times;
- 15           2.   improved lesional access and prolonged  
lesional-site retention (of up to several  
hours);
- 20           3.   essentially complete avoidance of uptake by  
normal liver, producing maximal lesional-to-  
normal organ enhancement ratios for hepatic  
tumors and other focal liver diseases;
- 25           4.   more rapid and complete body clearance, which  
occurs almost exclusively by the renal route;
- 30           5.   minimization of carrier side effects, which  
results from utilization of the lowest  
molecular weight fractions of heparin (e.g.,  
ca. 8,000 mean MW) commercially available in  
large quantity at acceptable cost;
- 35           6.   capability to use the heparin fractions and  
sources which have optimal sulfation ratios for  
both:
  - a.   multiple paired-ion binding of cationic  
MRI polyatomic complexes; and

-10-

- b. optimal complementarity to and interaction with the endothelial (and epithelial) determinants, including heparan sulfates, which appear to be responsible for lesional site selectivity.

7. optimal reformulation into extremely small, round, physically stabilized nanospheres (see below).

Also among the novel teachings set forth in the present application is reformulation of the preceding soluble MRI-carrier combinations in such a fashion as to provide new, extremely small nanospheres, typically less than the 100 nanometer diameter mean size previously disclosed by the applicant (PCT/US89/04295), and preferably about 5-99 nanometers in diameter. Such very small nanospheres confer, among others, the following major advantages:

1. the most rapid uptake of particles available from the blood plasma across endothelial (and epithelial) barriers into sequestered tissue sites and sites of disease, via "rapid" transport vesicles, at a rate which markedly exceeds that of larger particles with diameters greater than about 100-120 nanometers;
2. the most complete bioavailability possible of entrapped magnetic agents, against which water protons need to diffuse with minimal obstruction in order to afford maximal MRI relaxivity and in vivo potency per unit of entrapped magnetic species;



-11-

3. when desired (see below), optimal delayed dissociation of the entrapped diagnostic agents from the carrier substances, leading to rapid body clearance and maximizing the fraction of diagnostic agent which clears by the renal route.

Also among the novel teachings set forth in the present application is the reformulation of the preceding soluble MRI agent-carrier combinations, which incorporate low molecular weight heparin fractions of about 8,000 mean MW, into single-component and mixed-component, nanospheres of about 5-99 nanometers (mean diameter) as well as larger microspheres of about 0.1 to 250 um in mean diameter, to form:

1. agent-heparin microspheres with rapid bioavailability of entrapped MRI agent (or other actives), together with optimal spherical shape (in comparison to microspheres formulated with higher MW heparins in the mean MW range of about 16,000 to 22,000 Daltons).
2. mixed-component nanospheres and microspheres containing the magnetic agent entrapped in both heparin and protamine, hexadimethrine or starch matrix components, among others which range respectively and reciprocally from about 2 to 98 weight ratios, and wherein the second matrix component confers additional microsphere stability, as well as potentially (depending in part on diameter) extended or less commonly, accelerated release of the entrapped agent.

Also among the novel teachings of [those] set forth in the present application is that formulation of these

presently described nanoparticles and microparticles using the newly described lower-molecular weight heparin fractions, causes these particulate reformulations to have improved lesional-site and organ-site selectivity, and can confer other advantageous properties of these novel nanoparticles and microparticles within the animal or human being imaged.

Examples of optimal heparins include but are not limited to the more heavily sulfated, Fraction A heparins, and especially preferably those Fraction A heparins derived from beef-lung source. These heparins have not only improved sulfation ratios *in vitro*, but also improved complementarity to and interaction with endothelia (and epithelia) *in vivo*, as evidenced by improved *in vivo* binding to endothelium (and epithelium) and resultant site localization (see Examples). Such heparins classically include, and are herein further defined as including, various low molecular weight and rapidly eluting chromatographic fractions of heparins, such as those with a mean molecular weight of about 8,000 Daltons, those with a molecular weight range of about 6,000 to 10,000 Daltons, and fragments of those heparins which may range from about 1,000 Daltons to about 10,000 Daltons. Furthermore, such heparins may also be obtained from sources other than beef-lung, including but not limited to porcine mucosa, other animal species and organ sources, and from genetically engineered (recombinant DNA) prokaryotic or eukaryotic fermentation or cellular systems, among others. The heparins just described confer novel advantages, including *in vivo* site uptake and site selectivity for the soluble as well as the particulate, novel formulations described in the present application. Importantly, this subgroup of heparins provides such advantages even in the absence of the larger molecular size ranges of about 15,000 to 22,000

Daltons, previously thought to be necessary for prevention of free diffusion into normal tissues based on the mechanisms of size exclusion across capillary endothelial (and epithelial) barriers.

5

The size of both nanoparticles and microparticles are supramolecular. As taught in the present application, the optimal mean size of nanoparticles is less than about 100 nm in diameter, with the preferred mean diameter falling between about 5 and 99 nanometers. This is because "rapid" endothelial transport vesicles, whose activity is required for the very most rapid, selective transport of MRI enhancing agents from the blood (plasma) across microvascular endothelial barriers into the tissues, provide maximal transport rates and efficiencies for nanoparticles less than about 100 to 120 nanometers in diameter. Secondly, nanoparticles of less than about 100 nanometers in diameter allow the maximal quantity of entrapped magnetic agent to be made available rapidly to surrounding, diffusible magnetic atoms and molecules within body tissues. Thirdly, nanoparticles of this small size appear optimal for blood-pool image enhancement. It should be noted in this regard, that larger microparticles of 0.1 to 250 micrometers in diameter may also be formulated to undergo timely dissolution, such that essentially all of their entrapped magnetic material is also made available for interaction at pertinent distances of less than about 10 angstroms with surrounding diffusible magnetic species. Still, for formulations containing the same carrier (particle matrix) materials, the smaller nanoparticles typically dissolve more rapidly than do the larger microparticles.

Previous publications have disclosed entrapment of gadolinium chelates, such as gadolinium diethylenetriamine pentaacetic acid (Gd-DTPA), in

relatively small liposomes, in order to selectively enhance MRI images of the reticuloendothelial organs (liver, spleen and bone marrow) and potentially the lungs. (Buonocore et al., 2 Proc. Soc. Mag. Res. Med. 838 (1985) (which is hereby incorporated by reference).) Liver clearance is mediated by phagocytic (Kupffer) cells which spontaneously remove these small (0.05 to 0.1 micron) particles from the bloodstream (Buonocore et al. (1985) 2 Proc. Soc. Mag. Res. Med. 838). (Particles larger than 3 to 5 micron are selectively localized in the lungs, due to embolic entrapment in lung capillaries.) A recent report indicates that the small-sized Gd-liposomes produce effective decreases in liver T1's (as determined spectroscopically without imaging): see Buonocore et al. (1985) 2 Proc. Soc. Mag. Res. Med. 838). Also, insoluble Gd-DTPA colloids have recently been reported to enhance MR images of rabbit livers under *in vivo* conditions (Wolf et al. (1984) 4 Radiographics 66 (which is hereby incorporated by reference)). However, three major problems appear to limit the diagnostic utility of these devices. The multilamellar, lipid envelopes of liposomes appear to impede the free diffusion of water protons into the central, hydrophobic cores of these carriers, as assessed by the higher doses of Gd required for *in vitro* relaxivities equivalent to Gd-DTPA dimeglumine (Buonocore et al. (1985) 2 Proc. Soc. Mag. Res. Med. 838). This increases the relative toxicity of each Gd atom.

Even more importantly, these same lipid components cause the carriers to interact with cell membranes of the target organs in a way which leads to a marked prolongation of tissue retention, with clearance times of up to several months. (See Graybill et al., 145 J. Infect. Dis. 748 (1982) (which is hereby incorporated by reference), and Taylor et al., 125 Am. Rev. Resp. Dis.

-15-

610 (1982) (which is hereby incorporated by reference).) Two adverse consequences result. First, image enhancement does not return to baseline in a timely fashion. This precludes re-imaging at the short  
5 intervals (ca. 1 to 3-weeks) needed to assess acute disease progression and treatment effects. Second, significant quantities of the liposomally entrapped Gd-DTPA may be transferred directly into the membranes of host cells. (See Blank et al., 39 Health Physics 913  
10 (1980) (which is hereby incorporated by reference); Chan et al., 2 Proc. Soc. Mag. Res. Med. 846 (1985) (which is hereby incorporated by reference).) This can markedly increase the cellular retention and toxicity of such liposomal agents.

15

The consequences for Gd toxicity have not yet been reported. Protein (albumin) microspheres with entrapped Gd and Gd chelates have been prepared, and have been determined (by the present inventor and others: see Saini  
20 et al. (1985) 2 Proc. Soc. Mag. Res. Med. 896) to have only modest effects on T1 relaxivity *in vitro*. This is because most of the Gd as well as other entrapment materials are initially sequestered in the interior of these spheres, and are released very slowly as the  
25 spheres become hydrated (with  $t_{1/2}$ s of hours). (See Widder et al., 40 Cancer Res. 3512 (1980) (which is hereby incorporated by reference).) This phenomenon has been found by the present inventor to markedly reduce the acute (30-to-90-minute) relaxivity of each Gd atom to  
30 approximately 1/10th that of Gd-DTPA dimeglumine. Hence, both the quantity of carrier material and the toxicity of Gd are both unnecessarily high.

Emulsions of insoluble, gadolinium oxide particles  
35 have been injected into experimental animals, with significant image-enhancing effects on the liver.

(Burnett et al. (1985) 3 Magnetic Res. Imaging 65).  
However, these particles are considerably more toxic than  
any of the preceding materials, and are inappropriate for  
human use.

5

Novel Compositions and Methods for Imaging

Because of the significant disadvantages of existing  
MR image contrast agents, the present inventor has  
10 formulated improved agents with reduced toxicity,  
increased selectivity of tumor and organ uptake, as well  
as a significant potential for enhancing blood flow  
images.

15 A very important consideration, as taught by the  
present application, is that the marker substance should  
preferably be selectively deposited at the tissue  
location which is sought to be imaged. Moreover, the  
present application also contains significant teachings,  
20 regarding how the paramagnetic marker substance is bound  
to the polymer, which are believed to provide substantial  
advantages over previous teachings. The present  
application provides a novel method for NMR imaging,  
wherein image contrast is very strongly enhanced by a  
25 selective transport method which introduces an  
intramolecularly superparamagnetic strong T1 agent  
selectively into the desired imaging locations, and  
specifically into tumor locations. The present  
application also provides novel compositions of matter  
30 which are useful in implementing these methods.

MRI contrast enhancement can be improved moderately  
in the brain (and greatly in the body, which lacks the  
brain's tight blood-tissue barrier), by increasing the  
35 tumor selectivity of agent uptake.

-17-

Injected gadolinium exchanges off its DTPA chelator at a slow but significant rate *in vivo*. The resulting free gadolinium forms insoluble oxides, clears slowly from the body, and may produce significant side effects.

5 The present application permits major advantages to be gained, by substituting a less toxic, efficiently cleared, polyatomic metal-atom complex in an improved delivery process.

10 Chromium, in the form of  $^{51}\text{CrO}_4^{-2}$ , has been used extensively as a clinical agent for radionuclide labeling of platelets and red blood cells. Hexavalent chromate is converted within the red cell to the  $\text{Cr}^{+3}$  cation, which binds tightly but not irreversibly to hemoglobin.

15  $^{51}\text{Cr}$  elutes from red cells at an average rate of 0.93% per day. It is not reutilized by the body, but is cleared efficiently by excretory pathways. It is also of low toxicity in humans, even at relatively large doses. According to studies, in which neutrophils, tumor-cells  
20 and other biological targets have been labeled *in vitro*,  $^{51}\text{CrO}_4$  has been shown to bind to several cytoskeletal and cytoplasmic proteins (actomyosin as well as hemoglobin), and also to adenine nucleotides. It has also been shown to be nontoxic *in vitro* at relatively high concentration,  
25 as assessed by sensitive measures of cellular metabolism, DNA synthesis and cell division.

When tested as a potential MRI contrast agent, chromium (+3) has only moderate potency compared to gadolinium (+3), which has the highest number of unpaired  
30 electrons (7) of any metal ion. On this basis, the low-molecular-weight gadolinium chelate, Gd-DTPA, was developed as the first clinical MRI contrast agent, even though its small retained fraction (usually less than  
35 0.5%) is substantially more toxic than equivalent quant-

ities of retained chromium, and chromium is cleared much more completely than is gadolinium.

### Superparamagnetic Compounds

5

One class of highly paramagnetic compounds is those in which each molecule includes multiple atoms of magnetic moment with parallel spin vectors. While such intramolecular paramagnetic coupling does not imply that macroscopic ferromagnetic behavior will necessarily occur, it does imply that the resulting compound will be very strongly paramagnetic. Thus, such compounds are referred to as "intramolecularly superparamagnetic."

15

An article by Bino et al., "[Cr<sub>4</sub>S(O<sub>2</sub>CCH<sub>3</sub>)<sub>8</sub>(H<sub>2</sub>O)<sub>4</sub>]: Ferromagnetically Coupled Cr<sub>4</sub>S Cluster with Spin 6 Ground State", in the 9/16/88 issue of Science at page 1479, reported that the paramagnetic potency of chromium can be increased markedly by reformulating it as an intramolecularly ferromagnetically coupled cluster of four coordinated chromium ions, [Cr<sub>4</sub>S(O<sub>2</sub>CCH<sub>3</sub>)<sub>8</sub>(H<sub>2</sub>O)<sub>4</sub>]<sup>2+</sup>. This new, divalent chromium-organic complex cation has 12 unpaired electrons (1.7 times as many as gadolinium), and a magnetic spin of S = 6 (ground state). The effective magnetic moment of each Cr<sup>+3</sup> atom in the molecule is increased, due to stabilization of a coordination state which minimizes intramolecular antiferromagnetism.

25

The Bino et al. article refers to using the disclosed cation as a "spin label". (Spin labeling studies are normally *in vitro* studies.) The Bino et al. article also notes that the water ligands on the cation are potentially labile, so that the hydration sites would provide ligand bonding sites for association with potential carrier substances.

35



This tetra-chromium-sulfur-acetate complex is very advantageous for use in formulating MRI contrast agents of high potency and low toxicity. However, its small molecular size would cause it to equilibrate freely with  
5 the total extravascular (plasma + extracellular) water (as does Gd-DTPA), thereby reducing its potency and tumor/lesional-site selectivity.

#### Transport Properties of Polymeric Carrier

10

The present application teaches that major advantages can be gained by complexing or conjugating a superparamagnetic complex, such as  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , to polymeric or microspheric carriers which restrict its  
15 biodistribution, increase its selectivity of tumor localization, and amplify its proton relaxivity by slowing its rotational correlation time.

The present inventor has disclosed, in earlier  
20 filings, a method for increasing both the chemical potency (proton relaxivity) and tumor selectivity of paramagnetic contrast agents (including Gd-DTPA), by conjugating them to water-soluble, biocompatible carbohydrate polymers (including dextrans), whose  
25 molecular size distribution ranges from just above the cutoff for filtration out of normal microvessels (ca. 15,000 Daltons) to just below the cutoff for rapid renal clearance (ca. 45,000 Daltons). The present  
application discloses specific, improved, novel carriers  
30 and carrier-polyatomic cluster compositions which give improved site-selectivity even below the formerly taught 15,000 dalton cutoff.

-20-

Optimal Size Range

5 Selectivity of tumor uptake can be conferred in part  
by the polymeric size, in conjunction with characteristic  
changes in microvascular surface properties and an  
increase in porosity of malignant tumor microvessels.  
This allows the polymeric species (between 15,000 and  
10 45,000 Daltons) to filter or become transported more  
efficiently out of microvessels into the extravascular  
compartment (tumor interstitium). Due to the low  
porosity of normal microvessels, polymeric contrast  
agents are not allowed to filter into the surrounding  
normal tissues. This property of selective partitioning  
15 by molecular size: 1) advantageously results in steeper  
contrast gradients and increased intensity differences  
between tumor and normal tissues, and 2) advantageously  
produces highly discrete identification of tumor margins.  
However, there are major potential disadvantages to  
20 agents greater than 15,000 Daltons in size, as outlined  
above. Smaller agents, which are still site selective,  
represent a significant improvement.

Endothelial Binding

25 Tumor localization is also facilitated by the  
endothelial binding properties of negatively charged  
carbohydrates, especially highly sulfated, generally  
lower molecular weight heparins which are complementary  
30 to endothelial/epithelial determinants, including, but  
not limited to Fraction A heparins of about 6,000 to  
10,000 MW. (Simple polysaccharides will normally not  
adhere to the endothelial wall, unless the polysaccharide  
includes a charged surface group, such as sulfate,  
35 carboxyl, or dicarboxyl.)

-21-

thelium of normal tissues. Such prolonged binding at pathologic foci results in selectively accentuated uptake into the tumor interstitial gel proximal to sites of vascular endothelial binding.

5

#### Transendothelial Migration

Active endothelial transport has been demonstrated for small molecules (e.g., glucose and insulin).

10 However, no studies other than those of the present applicant are known to have shown such transport for larger molecules, or for molecules carried in a cargo format. It is now known (from the present applicant's histologic studies) that transendothelial migration of  
15 particles and molecular aggregates (larger than ca. 2 nm in diameter) can be accelerated by the application of appropriate surface coatings, preferably glycosaminoglycans or anionic polyglucoses or polyglycer-  
ols. (The glycosaminoglycans preferably include heparin,  
20 heparin derivatives and heparin fragments, but may also include dermatan sulfate, chondroitin sulfate, and other nature or modified glycosaminoglycans, including semisynthetic carboxylated glycosaminoglycans.) These surface coatings will bind multiply to receptors or  
25 antigens, which are either synthesized by endothelium or, although synthesized at other sites, become tightly associated with the endothelial surface. (See Ranney, 35 Biochem. Pharmacology 1063 (1986), which is hereby incorporated by reference). Such multiple binding  
30 typically involves complementary molecular interactions at more than 5 binding sites per molecule, and preferably more than 10 sites per molecule, and is termed adhesion, surface adhesion, or bioadhesion.

35 Following extravasation, the new agents in the present application are retained for prolonged intervals

in the tumor interstitium (greater than 2.5 hours, as compared to about 10-45 minutes for Gd-DTPA), and remain ("stay put") preferentially in the viable (versus necrotic) subregions. These two properties: 1) allow the agent to be injected at earlier times before imaging (and hence allowing premedication outside the imaging room); 2) permit tumor-treatment effects to be monitored in responding tumor subregions at early post-treatment intervals (at about 6-30 hours); and 3) allow viable and nonviable tumor to be distinguished at submillimeter resolution. This is because dead subregions cease to perfuse (and hence cease to take in the image-enhancing agents), while viable subregions continue to perfuse and take up image-enhancing agent. Partially damaged subregions continue to perfuse, and, since their microvessels typically have a still-further increased porosity (due to treatment-induced primary or secondary vascular damage), they allow the larger species of polydisperse polymer (as well as the smaller ones) to extravasate into the tumor gel. Hence, the partially damaged and potentially recoverable subregions achieve the brightest image intensity, because they accumulate the greatest quantity of these new, selective contrast-enhancing agents.

25

A further teaching of the present application is that these anionic glycosaminoglycans, polyglucoses and polyglycerols (and analogous compounds) undergo accentuated uptake by tumor cells, compared to the rate of uptake by normal cells in the same tissue region. This is based on the anionic (negatively charged) nature of side groups present on the polyglucose carriers, which engage the cellular uptake receptors of the anionic transport channels (pores) which are typically induced in hepatocellular carcinomas (hepatomas). Such anionic transport channels have also been found in several other

35

tumor types tested to date (by the present applicant and others). These same transport channels are relatively uninduced in the normal cell counterparts. This property of anionic small molecules and macromolecules facilitates  
5 active tumor-cell accumulation of the carrier polymer (and its bound ligands) *in vivo*. This property is exploited, in the innovative method disclosed herein, to allow for prolonged tumor retention and imaging.

10 It has previously been shown, by the present inventor as disclosed in a previous filing (International Application PCT/US88/01096), that IMFERON™ (which is a tightly bound, iron oxide-dextran complex of about 110,000 Daltons) achieves increased intramolecular  
15 paramagnetism (becomes superparamagnetic ) overall (e.g., intermolecularly as well). This complex is injected into patients for the purpose of achieving controlled iron release, over intervals of days to weeks, in order to treat iron deficiency anemia. Although it has been  
20 injected intravenously into patients, this must be done by controlled rather than bolus infusion, due to the release of a small fraction of its ionic iron which has been associated with acute toxicities. Hence, IMFERON™ is usually administered intramuscularly. These problems,  
25 together with the requirements, in MR image enhancement, of rapid intravenous administration of relatively large doses of the contrast agent, have precluded the effective use of IMFERON™ as an intravascular superparamagnetic contrast agent. However, this experimental evidence  
30 provides further confirmation that, as described below, metal coordinates of high potency and lower toxicity can be reformulated as polymeric agents (with a conjugation chemistry which is somewhat analogous to that of IMFERON™). Such metal-coordinate-polymer agents can be  
35 administered for purposes of tumor-selective MR image

enhancement, or alternatively to provide localized hysteresis superheating.

Use of Microaggregate or Microparticulate Carrier

5

In one class of embodiments, the carrier is used in the form of nanoparticles and microparticles. As discussed above, these particles have been found (when appropriately surfaced with sites complementary to endothelial determinants) to transport through the more porous parts of the endothelium walls with high preference. This is particularly advantageous in transporting a relatively high dose of the desired substance into the abnormal tissues and cells.

15

Such a nanosphere is most preferably between about 5 and 99 nanometers, and such a microsphere is most preferably between about 0.2 and 250 micron in diameter. The matrix of the microsphere is preferably a carbohydrate, and may be a carbohydrate such as heparin which also has multivalent binding capabilities especially a Fraction A heparin of about 6,000-10,000 Daltons or fragment thereof larger than about 1,000 Daltons. Protamine, hexadimethrine, starch and other matrix materials can also be used, and can also be coated with heparins. Such a microsphere carbohydrate matrix can optionally include, as a multivalent binding agent, an exposed or covert substance which is capable of binding endothelial surface determinants, enzymes, epien- dothelial or subendothelial substances. (Note that the nanoparticle/microparticle matrix may be coated with such a binding substance.)

In such embodiments, the nanoparticles/microparticles of the novel material disclosed herein will bind to endothelia (or to epithelia and

their closely associated extracellular structures), with preference (and longer residence times) in the vicinity of tumors (or other biological lesions if desired). This preferential binding leads to preferential induction, since a bound microsphere may be totally or partially enveloped in, for example, less than 10 to 15 minutes. The interaction of the preferred microspheres with endothelia may produce an induction of the endothelia to undergo transient separation or opening. The opening of the endothelia exposes underlying substances to which (ideally) binding may occur.

The present application provides improved methods and compositions of matter for the selective tumor localization of ferromagnetically coupled image-enhancing agents, contrast agents or spectral shift agents. This permits improved acquisition of tumor, tissue or organ images or spectra from live animals by nuclear magnetic resonance imaging or spectroscopy.

#### Additional Novel Compositions of Matter

It should be noted that the present application describes not only a number of novel methods, but also a number of novel compositions of matter, as set forth in greater detail below.

One novel teaching of the invention involves use of (I) a ferromagnetically coupled, multiply paramagnetic ion cluster (hereafter also designated the "intramolecularly superparamagnetic polyatomic complex") which is multiply associated, by complexation (including ion pairing) or covalent conjugation, with (II) a soluble, hydrophilic, biocompatible, excretable polymeric carrier, comprising repeating hydrophilic monomeric units, or with (III) monomeric or oligomeric subunits or

fragments of the final polymer, wherein the polymer or polymer subunits (either derived from natural sources or synthetic) have repeating monomeric units with a high frequency of hydroxyl, carbonyl, aldehyde, carboxyl, sulfate, sulfonate, sulfonium, phosphate, phosphonate, phosphonium, amine, amino, or quaternary ammonium groups, singly or in combination on the polymer, and the polymer contains less than about 5% (w/w) cross-linked or microaggregated species, all of low toxicity. The latter groups are for the purposes of either noncovalently binding the superparamagnetic complex or binding to target (including tumor) microvascular endothelium, or binding to both of the preceding entities.

The polymeric agent may optionally be formulated using an excipient counterion to achieve charge balance. Such excipient agents may include, for example, organic amines, preferably including N-methylglucamine (meglumine).

The superparamagnetic complex of the primary preferred embodiments uses a central tetrahedrally coordinated sulfur atom, surrounded by four octahedrally coordinated Chromium atoms, which are stabilized by bridging ligands, (which join pairs of Cr atoms). In the embodiment of Example 10, eight bridging ligands are used, and they are all acetate groups. However, in other embodiments, other bridging ligands, and/or a different number of bridging ligands, may be used.

The polymers most preferably used are Fraction A heparin (or heparan sulfate). However, of course, a large variety of other carrier polymers could be used instead. Note that the preferred polymer molecules are hydrophilic, which is required to provide the necessary environment for reliable NMR results.



### Alternative Carrier Compositions

Some of the other polymer species include other dextrans, dextran sulfate, dextran carboxylate, dermatan sulfate, chondroitin sulfate, pentosan polysulfate, hydroxyethyl starch, carboxylated hydroxyethyl starch or CARBETIMER™, carboxylated hydroxyethyl starch, and carboxylated dextrans in which the carboxylating groups consist essentially of multiple closely spaced carboxylates which are thereby capable of undergoing chelation-type or coordination-type binding with polyatomic organometallic complex structures which include metal ions.

The locations of the charged groups in the polymer can be readily modified, by methods well known to those skilled in the art, e.g., by introducing succinylate or glutaryl groups to extend the charge ion groups out from the polymer based structure. Thus, where it is desired to increase the affinity of the polyatomic unit being transported for the polymeric carrier molecule, the conformation of the polymer can optionally be modified in this fashion to achieve a better fit.

### Alternative Bridging Ligands in a Superparamagnetic Complex

The bridging ligands in the superparamagnetic complex need not be limited to acetate groups. A wide variety of organocarboxylates may be used. Some examples of alternative bridging ligands include: formate; formaldehyde; glutaraldehyde; glycinate; succinate; acetylacetonate; malonate; propionate; glutarate; hydroxamate; oxalate; 2-bromoacetate; 2-sulfoethanoate, thiolacetate; and thioglycolate.

### Use of Reactive Bridging Ligands

The embodiment described below, which includes at least some glycines as bridging ligands, has the advantage that the glycines contain sites which can assist in binding. Thus, a further secondary teaching is that the bridging ligand should contain a charged and/or activatable site.

### 10 Alternative Paramagnetic Species

The paramagnetic ion which is used in the superparamagnetic complex is most preferably chromium, but may alternatively be one or more of the following species: iron, nickel, manganese, cobalt, vanadium, molybdenum, tungsten, copper, platinum (particularly <sup>195</sup>Pt), erbium, gadolinium, europium, dysprosium and holmium.

### 20 Alternative Stabilizing Anions

The superparamagnetic complex  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8$  is preferably stabilized with water, so that the full formula of this complex cation is  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ . However, other stabilizing species can be used, such as sulfate, halide, nitrate, carboxylate, phosphates, or other stabilizing anions. (Note that some of these anions may be displaced when the complex binds to the endothelia or epithelia.)

### Additional Novel Methods

The present application also sets forth a generally applicable method for selective transport of a desired  
5 small polyatomic structure into tumors, or other regions of enhanced vascular porosity. Note that these novel teachings can be applied not only to the method of magnetic resonance imaging described, but also to a tremendous variety of other diagnostic and therapeutic  
10 uses.

### Further Points of Novelty

Among the novel teachings set forth in the present  
15 application is a method for magnetic resonance imaging, comprising the steps of: a) identifying a living vertebrate animal to be imagined; b) introducing into the blood stream of said animal a magnetic agent, comprising native or modified heparin or a fragment thereof, of  
20 molecular weight between about 1,000 and 10,000 Daltons, and a relatively small polyatomic structure which is intramolecularly superparamagnetic; c) applying to said animal a strong magnetic field which includes a gradient; d) and applying to at least a portion of said animal an  
25 electromagnetic perturbation field at a radio frequency generally corresponding to a resonant frequency of a predetermined species at a magnetic field strength which falls within the range of field strengths applied to said animal by said strong magnetic field, and measuring radio  
30 frequency response to define a spatial map of magnetic resonance characteristics within tissues of said animal, and e) analyzing said spatial map to determine the extent of tumors or other regions of enhanced vascular porosity.

35 Also among the novel teachings set forth in the present application is a method as above, wherein the

polymeric molecule has a molecular weight of about 10,000 Daltons or less.

5 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex comprises plural atoms of magnetic moment which are mutually ferromagnetically coupled.

10 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic structure is a polyatomic complex.

15 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex consists essentially of  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ .

20 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex comprises multiple atoms of magnetic moment selected from the group consisting of chromium, copper, nickel, manganese, platinum, erbium, gadolinium, erbium, dysprosium and holmium.

25 Also among the novel teachings set forth in the present application is a method as above, wherein said carrier is selected from the group consisting of: native or modified heparin, heparan sulfate, starch, protamine  
30 and hexadimethrine.

In this context, modified heparin, modified heparan sulfate and modified starch may include one or more of the reaction products of: acid hydrolysis, basic  
35 hydrolysis, hydrolytic preparation of lower molecular weight fragments than the native (parent) material or

purified fraction purified from parent material as obtained from its common source, enzymatic preparation of such fragments from native material or purified fraction of native material, derivitization with substituent  
5 groups which can modify association or binding of the polyatomic complex or of heparin, selected from: formate, formaldehyde, glutaraldehyde, acetate, glycinate, succinate, acetylacetonate, malonate, propionate, glutarate, hydroxamate, oxalate, 2-  
10 bromoacetate, 2-sulfoethanoate, thiolacetate, thioglycolate, carboxylate, ethylenediaminetetraacetate, diethylenetriaminepentaacetate, hydroxyethyl groups, carboxyethyl groups; reagents comprising multiple carboxylates, carbodiimide linking reagents, other  
15 linking reagents, chemical spacer groups, and other chemical derivitizations, alterations of physical conformation and partial degradations which comprise standard chemical and physical reactions performed on heparins, heparan sulfates, starch, simple and complex  
20 carbohydrates, and glycosaminoglycans, as well as the reactions (except for proteolysis) listed in the following paragraph.

In this context, modified protamine and  
25 hexadimethrine may include one or more of the reaction products of: combining these substances with one or more of native or modified heparin, heparan sulfate, and starch; crystallization, amorphous crystallization, and minor degradation which may occur upon one or more of  
30 partial proteolysis, exposure to organic solvents, exposure to mixtures of organic and aqueous solvents, heat, low pressure homogenization, high pressure homogenization, emulsification, combinations of the preceding reactions and processes, and other standard  
35 physical and chemical reactions performed on protamine, basic proteins, hexadimethrine and heparin antagonists.

Also among the novel teachings set forth in the present application is a method as above, wherein said carrier has a molecular weight of about 10,000 Daltons or less.

5

Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic structure comprises multiple metal ions mutually ferromagnetically coupled intramolecularly.

10

Also among the novel teachings set forth in the present application is a method as above, wherein all said constituents of said polyatomic complex, are reduced in standard measures of *in vivo* toxicity (relative to the free polyatomic complex) in animals and humans when the polyatomic complex is rendered in weak or strong chemical and weak or strong physical association with carrier (as defined above).

15

Also among the novel teachings set forth in the present application is a method as above, wherein said carrier comprises native or modified Fraction A heparin or a heparin fragment between about ,000 and 10,000 Daltons, which physically formulated as a microparticle having a diameter in the range of about 0.1 micron to 250 microns inclusive.

20

25

Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex is a cation which comprises multiple metal ions ferromagnetically coupled.

30

Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex is a cation which comprises

35

-33-

multiple metal ions complexed with multiple organic groups free of intermolecular bonds.

5 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex comprises metal ions, and wherein the toxicity of said metal ions is significantly reduced following association with said carrier.

10 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex is ion-pair bound to the interior of said carrier.

15 Also among the novel teachings set forth in the present application is a method as above, wherein said small polyatomic complex is covalently bound to the interior of said carrier by bonds which include at least one metal-oxide bond.

20 Also among the novel teachings set forth in the present application is a method as above, wherein a substantial fraction of said charged groups on the interior of said carrier are selected from the group  
25 consisting of: sulfides, amides, sites whose charge is strongly affected by an ether bond, and halides.

Also among the novel teachings set forth in the present application is a method as above, wherein said  
30 carrier has a molecular weight which is less than about 10,000 Daltons.

Also among the novel teachings set forth in the present application is a method as above, wherein said  
35 polymeric molecule has a mean molecular weight of about 8,000 Daltons.

Also among the novel teachings set forth in the present application is a method as above, wherein said carrier has anionic groups at the surface thereof.

5        Also among the novel teachings set forth in the present application is a method as above, wherein said perturbation field is applied at a frequency which generally corresponds to a resonance frequency of protons in an aqueous environment.

10        Also among the novel teachings set forth in the present application is an agent for image enhancement or spectral shift comprising: a strongly paramagnetic or intramolecularly superparamagnetic polyatomic complex  
15        having a spin of greater than  $3/2$ , more than about 7 unpaired electrons and labile or reactive ligands which can chemically or physically associate with: a biocompatible, excretable, water-soluble carrier comprising heparin having repeating hydrophilic monomeric  
20        units having hydroxyl, carboxylate, sulfate, or amine groups, singly or in combination on said carrier: wherein said image enhancing agent has a molecular weight of less than about 10,000 Daltons and is substantially completely water-soluble and contains less than about 5% (w/w)  
25        cross-linked or microaggregated species, all of low toxicity.

Also among the novel teachings set forth in the present application is an agent as above, comprising  
30        intramolecularly ferromagnetically coupled metal atoms of magnetic moment in association with the carrier, and wherein the image-enhancing agent is used to enhance internal images or shift internal spectra arising from induced magnetic resonance signals.



-35-

Also among the novel teachings set forth in the present application is an agent as above, wherein said carrier has a molecular weight between about 1,000 and 10,000 Daltons.

5

Also among the novel teachings set forth in the present application is an agent as above, wherein said intramolecularly ferromagnetically coupled polyatomic complexes are bound to said carrier noncovalently by a strong ionic (paired-ion or charge) interaction.

10

Also among the novel teachings set forth in the present application is an agent as above, wherein said carrier has a molecular weight between 1,000 and 10,000 Daltons, and said intramolecularly ferromagnetically coupled polyatomic complex is bound to said carrier by carboxylate or sulfate groups or both which are covalently conjugated to said polymer.

15

Also among the novel teachings set forth in the present application is an agent as above, wherein said atoms of magnetic moment are selected from said group consisting of chromium, copper, nickel, manganese, erbium, platinum, gadolinium, erbium, dysprosium and holmium.

20

25

Also among the novel teachings set forth in the present application is an agent as above, wherein said intramolecularly ferromagnetically coupled polyatomic complex is in a weight proportion to carrier of at least about 1:20.

30

Also among the novel teachings set forth in the present application is an agent as above, wherein said carrier is selected from the group consisting of: heparin, heparan sulfate, starch, hydroxyethylstarch,

35

carboxylated starch, carboxylated hydroxyethylstarch or carboxyethylstarch.

5 Also among the novel teachings set forth in the present application is an agent as above, wherein said polyatomic complex consists essentially of  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier consists essentially of heparin.

10 Also among the novel teachings set forth in the present application is an agent as above, wherein said polyatomic complex is  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  and said polymer is heparin.

15 Also among the novel teachings set forth in the present application is an agent as above, wherein said polyatomic complex is  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  and said polymer is DTPA-heparin.

20 Also among the novel teachings set forth in the present application is an agent as above, wherein said excess anionic charges of said carrier are balanced in part or totally by a nontoxic organic cation (base).

25 Also among the novel teachings set forth in the present application is an agent as above, wherein excess anionic charge of said carrier is balanced by N-methylglucamine (meglumine).

30 Also among the novel teachings set forth in the present application is an agent as above, wherein said polyatomic complex is covalently conjugated to said polymer totally or in part by a metal oxide linkage.

35 Also among the novel teachings set forth in the present application are the following:

1. an agent as above, wherein said intramolecularly ferromagnetically coupled atoms of the polyatomic complex comprise Cr(III), and said carrier comprise heparin having a molecular weight between about 5 6,000 and 10,00 Daltons, and having binding complementarity to determinants of mammalian endothelia and epithelia;
2. an agent as above, wherein said intramolecularly 10 ferromagnetically coupled atoms of the polyatomic complex comprise four Cr(III) atoms bound to a central tetrahedral sulfur atom and are octahedrally coordinated by bridging species, and said carrier comprises heparin having a molecular weight between 15 about 6,000 and 10,000 Daltons, and having binding complementarity to determinants of mammalian endothelia and epithelia;
3. an agent as above, wherein said polyatomic complex 20 comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  and said carrier comprises heparin having a molecular weight between about 1,000 and 22,000 Daltons, preferably 6,000 to 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia; 25
4. an agent as above, wherein said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprises Fraction A heparin having a molecular weight between about 6,000 and 10,000 Daltons and 30 binding to determinants of mammalian endothelia and epithelia;
5. In a preferred embodiment said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier 35 comprises Fraction A of beef-lung heparin having a molecular weight between about 6,000 and 10,000

Daltons, and complementarity to determinants of mammalian endothelia and epithelia.

- 5 6. In one aspect, said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprises native or modified Fraction A heparin or fragment thereof, having a molecular weight between about 1,000 and 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia.
- 10 7. In another embodiment wherein said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprises heparin having a mean molecular weight of about 6,000 Daltons, and binding determinants of mammalian endothelia and epithelia.
- 15 8. In another aspect said polyatomic complex is in association with said carrier noncovalently through a strong ionic, paired-ion or charge interaction, said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprises heparin having a molecular weight between about 6,000 and 10,000 Daltons, and having binding complementarity to determinants of mammalian endothelia and epithelia.
- 20 9. Said polyatomic complex may also be in association with said carrier noncovalently through a strong ionic, paired-ion or charge interaction, said strong interaction involves chemical coordination or chelation binding of said polyatomic complex to at least one reactive group of said carrier, each reactive group on said carrier having a coordination number or ionic charge number between 2 and 10.
- 25 30 Said polyatomic complex may comprise  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprise
- 35

heparin having a molecular weight between about 6,000 and 10,000 Daltons, and having binding complementarity to determinants of mammalian endothelia and epithelia.

5

10. Said polyatomic complex may also be in association with said carrier covalently, and wherein said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , and said carrier comprises heparin a molecular weight between about 6,000 and 10,000 Daltons, and having binding to determinants of mammalian endothelia and epithelia.
11. In one embodiment the association of the polyatomic complex with the carrier is stabilized by heating a pre-dried polyatomic complex plus carrier combination to at least about 100 degrees Celsius in one or more of an organic solvent, and an organic solvent comprising a one or more of a detergent or surfactant, a polyol, a biocompatible oil; and wherein said carrier comprises heparin having a molecular weight between about 6,000 and 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia.
12. In certain cases, association of the polyatomic complex with the carrier is stabilized by heating a pre-dried polyatomic complex plus carrier combination to at least about 100 degrees Celsius in an organic solvent comprising one or more of: acetone, acetone with polyoxyethylene sorbitan mono-oleate between about 0.01 and about 25 weight percent, polyethylene glycol, glycerol, soybean oil, and corn oil.

35

13. When said polyatomic complex,  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , association of the polyatomic complex with the carrier may be stabilized by heating a pre-dried polyatomic complex plus carrier combination to at least about 100 degrees Celsius in an organic solvent. Pre-drying of the polyatomic complex and use of an organic solvent during heating are to protect said  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  against chemical degradation which can occur slowly for free (carrier-unassociated)  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  in aqueous solvents and which can be accelerated at temperatures substantially above 22 degrees Celsius, most particularly at temperatures above about 60 to 80 degrees Celsius.
14. When said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , said carrier may comprise heparin having a molecular weight between about 6,000 and 120,000 Daltons, and binding to determinants of mammalian endothelia and epithelia, and, further, the heparin may be in combination with a polyamine such as protamine and hexadimethrine, for example in a reciprocal weight ratio of between about 98:2 and about 2:98;
15. Said intramolecularly ferromagnetically coupled polyatomic complex of  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  may be combined with a carrier comprising Fraction A of beef-lung heparin having a molecular weight between about 6,000 and 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia, and further the heparin may be in combination with one of: starch, hydroxyethylstarch, carboxylated starch, hydroxyethylstarch, carboxyethylstarch, ethylenediaminetetraacetate starch, diethylenetriaminepentaacetate starch, and other

modified starch, in a reciprocal weight ratio of between about 98:2 and about 2:98.

16. Said polyatomic complex may also comprise  
5  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ , said carrier comprising Fraction A of beef-lung heparin having a molecular weight between about 6,000 and 10,000 Daltons, binding to determinants of mammalian endothelia and epithelia, and wherein the heparin is in combination  
10 with one of: dextran, ethylenediaminetetraacetate starch, diethylenetriaminepentaacetate dextran, carboxylated dextran, dextran sulfate, and other modified dextran, in a reciprocal weight ratio of between about 98:2 and about 2:98.
17. Said polyatomic complex of  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  may  
15 be combined with a carrier comprising heparin having a molecular weight between about 1,000 and 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia.
18. Said intramolecularly ferromagnetically coupled  
20 polyatomic complex of  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$  may be combined with a carrier comprising ethylenediaminetetraacetate-heparin and  
25 diethylenetriaminepentaacetate-heparin, the carrier having a molecular weight between about 1,000 and 120,000 Daltons, and binding to determinants of mammalian endothelia and epithelia.
19. Said carrier may comprise heparin between about  
30 1,000 and 10,000 Daltons and be in association with said polyatomic complex, and wherein said polyatomic complex and said carrier are further formulated to  
35 compose nanoparticles stabilized by one or more of: chemical binding of two or more carrier substances,

heat, homogenization, physical dispersion, and chemical treatment. The nanoparticles have a mean diameter of between about 5 and 99 nanometers.

- 5    20. When said carrier comprises Fraction A of beef-lung  
heparin between about 6,000 and 10,000 Daltons and  
is in association with said polyatomic complex, said  
polyatomic complex and said carrier may be further  
10    formulated to compose microparticles, which are  
stabilized by one or more of: chemical binding of  
two or more carrier substances, heat,  
homogenization, physical dispersion, and chemical  
treatment. The microparticles have a mean diameter  
of between about 0.1 and 250 micrometers.
- 15    21. When said carrier comprises Fraction A of beef-lung  
heparin between about 6,000 and 10,000 Daltons and  
is in association with said associated polyatomic  
complex, said polyatomic complex and said carrier  
20    may be further formulated by combination with at  
least one of protamine, hexadimethrine, starch, and  
dextran comprised in said carrier, to compose  
microparticles. There may be stabilized by one or  
25    more of: chemical binding of two or more carrier  
substances, heat, homogenization, physical  
dispersion, and chemical treatment, and wherein the  
microparticles have a mean diameter of between about  
0.1 and 250 micrometers.
- 30    22. The proportion of said polyatomic complex to said  
carrier is at least about 1:20 by weight, in order  
that sufficient agent can be injected for MRI image  
enhancement (or spectral shift) without overloading  
the recipient animal with carrier material.



23. When the associated agent plus carrier, prior to any reformulation as nanoparticles or microparticles, preferably contains less than about 5 weight percent cross-linked or microaggregated species, in order to optimize pharmacokinetics (plasma circulation time), biodistribution and body clearance.

The present invention will be described with reference to the accompanying drawings, which show important sample embodiments of the invention and which are incorporated in the specification hereof by reference, wherein:

Figure 1 shows a perspective view of a sample superparamagnetic complex which can be used in one embodiment of the disclosed method.

Figure 2 schematically shows a nuclear magnetic resonance imaging system suitable for use in the disclosed innovative method.

Figure 3 is a schematic representation of the transport of microspheres entirely across endothelial (or epithelial) tissue.

Figure 4 is a photograph of stained tissue showing localization of nanospheres. The nanospheres, at the scale of this photograph, appear as small gray round or oval dots of 1 to 2  $\mu\text{m}$  in diameter.

Figures 5A-5E show infrared data pertinent to the fabrication of a further alternative superparamagnetic complex, which can be used instead of that shown in Figure 1. Figure 5A shows an infrared spectrum trace for the reaction products where  $\text{Cr}_2\text{S}(\text{O}_2\text{CCH}_3)_4$  was heated with glycine in water at  $92^\circ\text{C}$ . Figure 5B shows a trace for the reaction products where  $\text{Cr}(\text{NO}_3)_3$  was heated with glycine in water at

92°C. Figure 5C shows a trace for the reaction products where  $\text{Cr}_4\text{S}(\text{Ac})_8$  was refluxed in acetic anhydride for several hours. Figure 5D shows a trace for  $\text{Cr}_4\text{S}(\text{Ac})_8$  alone (which has a blue or green color), and Figure 5E shows a trace for  
5 glycine alone.

Figures 6a, 6b, 6c, and 6d show the following, representative infrared spectra. Figure 6a shows native chromium-ion cluster (with characteristic absorption maxima  
10 at 650-700 ( $\text{cm}^{-1}$ ) and 3200 ( $\text{cm}^{-1}$ ) wave numbers; Figure 6b shows formulation 009-59A (see also Table 5), containing chromium-ion complex formulated with pentosan polysulfate, and lyophilized (NOTE: characteristic sharp absorbance at  
15 650-700 ( $\text{cm}^{-1}$ ) and characteristic shoulder at 3200 ( $\text{cm}^{-1}$ ) of the native chromium-ion complex); Figure 6c shows formulation 009059B (see also Table 5), containing chromium-ion complex formulated with pentosan polysulfate plus protamine (protamine sulfate) as a minor component, and lyophilized (NOTE: characteristic sharp absorbance at  
20 650-700 ( $\text{cm}^{-1}$ ) and characteristic shoulder at 3200 ( $\text{cm}^{-1}$ ) of the native chromium-ion complex); Figure 6d shows pentosan polysulfate -- as a control for Figures 6a-c (above).

25 The numerous innovative teachings of the present application will be described with particular reference to the presently preferred embodiment, wherein these innovative teachings are advantageously applied to the particular problems of MRI imaging by selective  
30 localization of superparamagnetic molecules. However, it should be understood that this embodiment is only one example of the many advantageous uses of the innovative teachings herein. For example, the various types of innovative methods and compositions disclosed herein can  
35 optionally be used to selectively localize radionuclide, x-ray contrast and ultrasound/acoustic image enhancers,

-45-

or therapeutic agents. In general, statements made in the specification of the present application do not necessarily delimit any of the various claimed inventions. Moreover, some statements may apply to some  
5 inventive features but not to others.

Figure 2 shows an example of a nuclear magnetic resonance imaging (MRI) unit, useable for whole-body imaging of human patients. A patient 240 is inserted, on  
10 a sliding table 230, into the interior of a large solenoidal winding 210. The large coils 210 (which may be water-cooled or superconducting) will apply a constant (DC) magnetic field, typically 0.5 to 1 Tesla. (This field component is referred to as the  $B_0$  field.) Bias  
15 coils 220 apply a gradient to this field, as described above. Finally, a probe coil (which is movable, and is not shown in this Figure) is used to apply the RF pulses described above. Differently shaped probe coils are used for imaging different parts of the body, and the probe  
20 coil is often shaped so that it will nearly fit to the shape of the surface of the area to be imaged.

The present invention provides an improved MRI imaging method, whereby the ability of MRI systems to  
25 detect tumors is greatly enhanced. This is accomplished by selectively introducing an image-enhancing, spectral shift responsive agent into the abnormal tissue.

One novel teaching of the invention involves use of  
30 (I) a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are individually bound to at least one of said plural atoms  
35 having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of

intermolecular bonds, more preferably wherein said plural atoms having magnetic moments comprise chromium (III) atoms, more preferably wherein the polyatomic complex comprises four chromium (III) atoms bound to a central tetrahedral sulfur atom and the chromium (III) atoms are octahedrally coordinated by bridging species, and most preferably wherein said polyatomic complex comprises  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]^{2+}$ ; wherein multiple ones of polyatomic complex are associated by one or more of physical means, chemically by noncovalent binding, hydrogen bonding, complexation, strong ionic binding (charge interaction or paired-ion binding), coordination, or covalent conjugation; with (II) a soluble, hydrophilic, biocompatible, excretable, polymeric or oligomeric carrier, preferably comprising heparin in the molecular weight range of about 1,000 to about 10,000 Daltons, and more preferably, rapidly chromatographing Fraction A heparin in the molecular weight range of about 6,000 to about 10,000 Daltons, and most preferably Fraction A heparin in the molecular weight range of about 6,000 to about 10,000 Daltons having a mean molecular weight of about 8,000 Daltons and derived from beef lung; with (III) a fragment or oligomer prepared from (II, above); with (IV) a nanoparticle physical formulation prepared at least in part from (II or III, above), wherein the nanoparticle size is between about 5 and 99 nanometers in mean diameter; with (V) a microparticle physical formulation, prepared at least in part from (II, above) and preferably between about 0.1 and 250 micrometers mean diameter, and most preferably between about 0.1 and 15 micrometers in mean diameter, wherein said heparin is more preferably rapidly chromatographing Fraction A heparin in the molecular weight range of about 6,000 to about 10,000 daltons, and is most preferably beef-lung Fraction A heparin in the molecular weight range of about 6,000 to about 10,000 Daltons having a mean molecular

weight of about 8.000 Daltons; wherein the polymeric carrier (II or III, above) is derived from natural, synthetic or recombinant genetic sources and has repeating monomeric units with a high frequency (by  
5 natural or synthetic means) of hydroxyl, carboxyl, aldehyde, sulfate, sulfonate, sulfonium, phosphate, phosphonate, phosphonium, amine, amino, secondary ammonium or quaternary ammonium groups, singly or in combination on the carrier, said groups being for one or  
10 more of: association of said polyatomic complex (I, above) with said carrier substances (II or III, above) and binding said polyatomic complex-carrier agent to microvascular endothelium (or epithelium) including the endothelium of tumors and other lesional sites (sites of  
15 disease); wherein the weight ratio of said polyatomic complex (I, above) to said carrier (II or III, above) is at least about 1:20; wherein said carrier (II or III, above) is of low mammalian toxicity and contains less than about 5 weight percent cross-linked or  
20 microaggregated species; and wherein the carrier may comprise (VI) one or more additional substances in order to provide optimal stability, optimal availability of said atoms of magnetic moment, optimal *in vivo* properties, (VI) being one or more of native or modified  
25 starch, and native or modified dextran, protamine and hexadimethrine, at a weight ratio to heparin of between about 98:2 and 2:98; and (IV) being most preferably protamine. Said atoms of magnetic moment comprise one or more of: chromium, copper, manganese, iron, platinum  
30 (particularly  $^{195}\text{Pt}$ ), cobalt, vanadium, molybdenum, tungsten, gadolinium, erbium, dysprosium, europium and holmium, and preferably comprise chromium (III). Bridging species comprise one or more of: formate, formaldehyde, glutaraldehyde, acetate, glycinate,  
35 succinate, acetylacetonate, malonate, propionate, glutamate, hydroxamate, oxalate, 2-bromoacetate, 2-

5 sulfoethanoate, thiolacetate, thioglycolate, or the like,  
and preferably comprise acetate. Additionally,  
counterions may be used to stabilize the polyatomic  
complex in various aqueous and nonaqueous solvents and to  
10 optimize association with said carrier., Such  
stabilizing counterions may comprise one or more of:  
halide, sulfate, nitrate, carboxylate and phosphate, or  
the like, and preferably comprise chloride or sulfate.  
In certain cases and preferably for nanoparticle and  
15 microparticle formulations, the association of said  
polyatomic complex (I, above) and carrier substances (II  
through VI, above) may be further stabilized by physical  
means, including (VII) heating a pre-dried polyatomic  
complex plus carrier substance combination to at least  
20 about 100 degrees Celsius in an organic solvent, and in  
the absence or presence of homogenization, physical  
dispersion, or emulsification. Said organic solvent  
preferably comprises one or more of: acetone, a polyol,  
polyethylene glycol, glycerol, a detergent, a surfactant  
25 and a biocompatible oil; more preferably comprises  
acetone and a biocompatible detergent; and most  
preferably comprises acetone and polyoxyethylene sorbitan  
mono-oleate at between about 0.01 weight percent and  
about 25 weight percent. Said agent comprising said  
30 polyatomic complex (I, above) and one or more carrier  
substances (II through VI, above) and stabilized  
according (VII, above), may be further formulated using  
various excipients or vehicles, including a counterion in  
order to achieve charge balance. Such counterions may  
35 include inorganic or organic amines, preferably including  
the organic amine, N-methylglucamine (meglumine). The  
use of such excipients or vehicles may be either for the  
purposes of in vitro formulation and processing, or for  
the purposes of improving in vivo properties.

Additional preferred polymer substances include heparin, heparan sulfate, dextrans, dextran sulfate, dextran carboxylate, dermatan sulfate, chondroitin sulfate, pentosan polysulfate, hydroxyethyl starch, 5 carboxylated hydroxyethyl starch or CARBETIMER™, and especially heparin, carboxylated hydroxyethyl starch and carboxylated dextrans in which the carboxylating groups consist essentially of multiple closely spaced carboxylates which are thereby capable of undergoing chelation- 10 type or coordination-type binding with said polyatomic complex structures which include atoms of magnetic moment.

The ferromagnetically coupled polyatomic complex 15 includes a molecular coordination compound containing a paramagnetic metal atom present in numbers of two or more per molecular coordinate, wherein the paramagnetic metal ion includes one or more of the following: chromium, iron, nickel, manganese, cobalt, vanadium, molybdenum, 20 tungsten, copper, platinum (particularly <sup>195</sup>Pt), erbium, gadolinium, europium, dysprosium or holmium; and the complexed metal atoms are stabilized in a ferromagnetic or superparamagnetic intramolecular complex configuration by molecular or atomic bridging species. The preferred 25 metal atoms include chromium and gadolinium, and the preferred bridging species include organosulfates and their derivatives, carboxylic acids, and especially acetate ions. The polyatomic complex may also include a central multivalent stabilizing ion or element, in which 30 case the preferred element includes sulfur. The intramolecularly ferromagnetically coupled, polyatomic complex preferably has a net nuclear spin of greater than about 3/2 and has more than about 7 unpaired electrons. It has one or more of labile water groups a net 35 electrical charge, or reactive groups or combining sites

which allow it to chemically associate, covalently or noncovalently with the carrier polymer.

5 In operation, time-domain windowing may be performed in the MIR imaging run, so that the atoms with the shortest relaxation times are seen preferentially. Alternatively, a high pulse repetition rate may be used, so that the tissues with the longest relaxation times are kept in saturation.

10

One preferred group of features of the complex for chemical association includes carboxylate, oxygen, metal (especially chromium "hydroxide") glycine amine, and net cationic (positive) charge.

15

20 The heparin polyatomic complexes just described, are preferred for selective uptake by systemic tumors, liver and body tumors, lung, lung tumors and other lung lesions, for MRI image enhancement, or hysteresis hyperthermia, following intravenous injection, or particularly for said microparticle formulations, for uptake by solid tumors and other focal disease following selective arterial administration. Lung uptake can be increased by leaving a sufficient fraction of heparin's sulfate groups, preferably greater than 30%, unbalanced by counterions. Lung uptake can be reduced and systemic access increased by balancing more completely the negative sulfate groups of heparin with either or both of the superparamagnetic complex or the excipient counter-  
30 ion, including N-methylglucamine.

In certain instances, it may be preferable to separate the polyatomic complex from the polymeric carrier, using a linker which has a chain length of preferably between about 4 and 8 carbon atoms. This may  
35 be desirable under circumstances in which a) the



polyatomic complex is a bulky molecule, or b) it is important to stabilize the bond between the polymer and the linker against lysis, including hydrolysis and esterolysis, which may in unusual cases, be catalyzed by  
5 a substituent of the polyatomic complex.

In an alternative embodiment, a method of modifying the spatial relation of the magnetic substance to carboxylated carbohydrate polymers, is specified. This  
10 alternative method involves derivatizing the carbohydrate polymer with a higher bifunctional acid, including preferably succinic acid, to form a succinylated polymer with a 4-carbon spacer between the polymer and the polyatomic complex to be conjugated to the polymer via  
15 the free carboxylate group of each succinate linker.

Although the preceding preferred methods of conjugation have focused on ester and carboxylate linkages, other linkers are not excluded, and in some  
20 cases may be preferred. These include aldehyde, amine, amide, carbodiimide, halogen-activated carbohydrate groups, and combinations thereof.

A third preferred embodiment involves providing a  
25 physical microsphere or nanosphere form of the preceding agents, in which the diameter of the spheres ranges from 0.1 to 250 micrometers, and the particles are preferably formed from their paired, carrier-polyatomic complexes or from simple mixtures of the polymer and the substance to  
30 be entrapped. Matrix polymers and excipients preferably comprise between about 5% and about 75% of the particle weight. The particles are prepared preferably by phase dispersion emulsification (for larger ones) or high-pressure homogenization (for smaller ones), followed by  
35 heat or chemical stabilization of the polymer matrix, and extraction of the oil phase with an organic solvent,

including acetone, ether, or hexane, preferably acetones which may also contain a small quantity of a biocompatible detergent for surface stabilization, preferably polyoxyethylene sorbitan mono-oleate (or deoxycholate) at about 0.01 to 25% (w/w). Smaller particles are provided by high pressure homogenization. The degree of heat or chemical stabilization will preferably determine how long the particle retains its physical form following rehydration for *in vivo* administration, and will also determine how rapidly the internally entrapped substance of magnetic moment is made available, hydrated, exposed or released, in order that it can modify the biochemical environment of the plasma, extracellular matrix (or matrix water), or intracellular cytoplasmic constituents (or water). Preferably heat stabilization of carbohydrate matrices is performed for about 30 seconds to 5 minutes, in order to render the matrix sufficiently stabilized that the entrapped material becomes chemically exposed over an interval of about 15 minutes to 30 hours. For both the induction of MRI contrast and the amplification of hysteresis heating, the  $t_d$  for release of entrapped substance of magnetic moment occurs preferably within about 15-20 minutes of injection, although under certain circumstances, particularly those involving the monitoring of controlled-release drugs from selectively localized microcarriers, this  $t_d$  may be considerably longer. Also, for MRI contrast enhancement, the entrapped magnetic material, if paramagnetic, must be hydrated (released) in order to affect surrounding diffusible water protons, whereas, for hysteresis heating, the entrapped material need not be released at all, but can function while still entrapped, providing that the macrodomain size of an average superparamagnetic or ferromagnetic deposit within the particle is sufficiently large for efficient hysteresis to occur, preferably larger than about 0.5 micron. Otherwise,

-53-

release of entrapped magnetic material with subsequent reconcentration by extracellular matrix binding or cellular processes, is preferred in order to achieve an efficient hysteresis response in the tissues.

5

The smaller particle sizes (of less than about 10 to 15 micrometers, and especially less than about 3 micron) are preferred for systemic administration by intravenous injection and for selective arterial administration into critical end-arterial circulations. The larger particle sizes of greater than 15 micrometers, and especially greater than 100 micrometers are preferred for chemoembolization of selected organs with blood supplies, including especially the liver, by selective arterial administration, and for introduction mechanically, directly into tumor masses or body cavities.

The acute enhancement of blood flow (or perfusion) images, for example in the heart or cerebral vessels, may be accomplished with the soluble polymeric image-enhancing agents and is even more efficiently performed with the nanosphere and microsphere forms.

A significant advantage of MRI enhancement with polymeric, nanoparticle and microparticle formulations, is a further reduction of the dose and any potential toxicity over that which can be achieved by simple (low molecular weight) paramagnetic substances alone.

The relatively rapid biodegradation and metal clearance times, and the resultant shorter reimaging intervals are particular advantages involved with the present invention relative to other polymeric and particulate metal oxides, chelates and complexes.

35

The image-enhancing agents of the present invention, in soluble nanoparticle or microparticle form, are readily reconstituted for animal and patient administration. This reconstitution involves a simple  
5 vortex-type mixing, as contrasted with the sonication in detergents used for protein-based microspheres.

The image-enhancing agents of the present invention are easily usable in any MRI detection system involving  
10 administration of paramagnetic, superparamagnetic or ferromagnetic contrast agents. It has particular advantages in conjunction with the newer rapid RF pulse sequences, which reduce native tissue contrast in order to shorten image acquisition times and increase patient  
15 throughput. The image or spectral enhancing agents of the present invention allow shorter image acquisition times for satisfactory internal resolutions. These times are generally adequate to produce satisfactory internal images because of the greater enhancement and image  
20 contrast produced per unit of superparamagnetic and total agent.

The potential for selective localization of large numbers of relatively nontoxic intramolecularly super-  
25 paramagnetic molecules by small numbers of monoclonal antibodies, nonpeptide and peptide hormones, lymphokines, cytokines, and other receptor-binding substances tagged with one or more of the innovative (and preferably polymeric) image-enhancing agents is contemplated as a  
30 major diagnostic advancement for future use.

The potential for selective localization of large numbers of relatively nontoxic intramolecularly super-  
paramagnetic molecules by small numbers of carrier  
35 polymers, nanoparticles or microsparticles is contemplated as a major therapeutic advantage for future use in

conjunction with hyperthermia augmentation by hysteresis heating and delivery and monitoring of tagged therapeutic agent localization in sites of disease.

5       Because of the high MRI contrast conferred by these intramolecularly superparamagnetic substances and the substantial prolongation of lesional residence times, use of the present image-enhancing agents will allow an increased number of serial images to be obtained in an  
10       enhancement mode after a single administration of agent.

Due to the selective retention of the carriers used to formulate the present image-enhancing agents, superior definition of tumor margins and markedly improved  
15       discrimination of viable and nonviable tumor subregions is possible. This has the major advantage of allowing tumor responses to chemotherapy and radiation therapy to be monitored at early posttreatment times and submillimeter resolution, several weeks before small  
20       tumor nodules would regrow to volumes detectible by computerized axial tomography (CAT) and radionuclide scanning.

From a chemical standpoint, some advantages of the  
25       present invention may be summarized as follows. When MRI image-enhancing agents comprise intramolecular superparamagnetics, each magnetic substance exhibits an increased (strong) paramagnetic relaxivity for adjacent magnetic nuclei (e.g., protons) and hence, gives greater  
30       T1-weighted signal enhancement. This increased relaxivity is related to an increased dipolar correlation time of the superparamagnetic substance due to its slower molecular rotation when polymerically controlled. Spacer groups are not required between the intramolecularly  
35       superparamagnetic substance and the polymeric carrier in order to obtain optimal paramagnetic relaxation

potencies, however, they could be introduced if deemed advantageous for other purposes.

5       The chemically defined nature of preferred polyatomic complex-carrier combinations allows ready batch-to-batch uniformity for improved pharmaceutical formulations and a likely greater ease of regulatory approval.

10       Many of the preferred polymers of the present invention, such as certain dextrans (40,000 and 70,000 MW forms), hydroxyethyl starch, and heparins, for example, have already separately achieved final regulatory  
15       approval for patient administration. The size of these polymers is optimized to prevent access into normal tissues, but to still allow rapid renal clearance and essentially complete body clearance. This is especially true for the present novel heparins of less than about 10,000 Daltons. Also, due to the association of  
20       multiple, intramolecularly superparamagnetic substances with each polymeric carrier molecule, the resulting complexes and conjugates comprise low osmolality agents by comparison to their low-molecular weight counterparts. Such low osmolality agents have been shown  
25       to have major advantages in several categories of high-risk (particularly cardiovascular) patients.

For parenteral administration, these agents are preferably formulated as a sterile, physiologically  
30       balanced, aqueous solution (or suspension), whose pH for purposes of intravenous administration is 6.5 to 7.0. Alternatively, these agents may be lyophilized or spray dried and provided in the dried form for reconstitution in physiologic solutions just prior to administration.  
35       For injection into body cavities (such as the bladder, uterus, Fallopian tubes, nasal sinuses or ventriculo-

-57-

cerebrospinal system), these agent may be formulated as a physiological solution (or suspension) which contains additional substances (excipients) to increase the viscosity or osmolality. Other additives and  
5 formulations may also be incorporated according to standard pharmaceutical procedures.

For parenteral administration, the concentration of total active agent (polymer-superparamagnetic substance)  
10 should be between about 0.1% and 30% (weight/volume), typically between about 5% and 25%, and preferably about 20%. Doses of the soluble polymer, nanoparticle and microparticle agents will vary depending on the super-  
paramagnetic substance used and the route of  
15 administration. The following doses are given for intravenous administration. For tumor image enhancement with some of the preferred embodiments, which include soluble  $\text{Cr}_2\text{S}(\text{O}_2\text{CCH}_3)_2(\text{H}_2\text{O})_4^{+2}$  complexed to heparins, the dose of chromium will be between about 0.005 and 0.01  
20 millimoles per kilogram of body weight. For enhancement of the cardiovascular blood pool, the optimal dose will occur at or below about 0.04 and 0.3 millimoles of chromium per kilogram.

25 For hysteresis heating, the nanosphere or microsphere forms of agents will be administered once or multiply, at about 15 minutes to 2 hours prior to each treatment by either systemic intravenous injection, direct administration into superficial tumors, or by  
30 intraarterial perfusion, at doses of up to 1.0 mmol/kg of the superparamagnetic substance. Hysteresis hyperthermia at a frequency of about 10 to 150 kHz will be directed from an external oscillating magnetic source whose maximal energy displacement is centered over the major  
35 external or imageable mass(es) of tumor. A major anticipated advantage of using the nanosphere or microsphere

form of preferred embodiment, is that a high concentration of clustered (macrodomain) metal is selectively localized in the diseased subregions of the target tissue following transvascular administration, and this  
5 localized material further concentrates in the viable, most heavily perfused subregions of tumor which require the greatest augmentation of heating to compensate for their disproportionate loss of heat due to blood flow dissipation.

10

Figure 3 schematically shows how the trans-epithelial transport of microspheres works. A small blood vessel 310 is shown passing through tissue 320. Four microspheres (or nanospheres) 301, 302, 303, and 304  
15 are shown at different stages of passage.

20

Microsphere 301 has recently adhered to the endothelial wall 311. Thus, this microsphere is said to be at the stage of endothelial adhesion.

25

Microsphere 302 is at the stage of endothelial envelopment. This will occur after a few minutes, as the wall 311 gradually covers a particle (like microsphere 301) which has adhered to it.

30

Microsphere 303 is shown at the further stage of extravasation, after envelopment has passed it entirely across the endothelial (or epithelial) wall 311.

Finally, microsphere 304 is shown at yet a further stage, of percolation through tissue.

35

Figure 4 is a photograph of stained tissue showing lung localization of heparin-coated subembolizing (0.1 - 0.8 micron) nanospheres distributed throughout rodent lung tissue at 5 minutes after intravenous injection via



the tail vein. The nanospheres, at the scale of this photograph, appear as small gray round or oval dots of 1 to 2 mm in diameter.

- 5 Additional novel approaches and formulations described herein include but are not limited to the use of acidic saccharides which include phosphated saccharides, carboxylated saccharides sulfated saccharides and heparin (which itself contains both
- 10 carboxylate and sulfate groups). Typical sulfated saccharides include pentosan polysulfate and sucrose octasulfate, which have higher numbers of sulfate substituents than heparin. They therefore have a higher binding capacity per saccharide unit than does heparin.
- 15 Whereas, heparins typically have about 2.8 to 3 sulfate groups per disaccharide unit, pentosan polysulfate has about 4 sulfate groups per disaccharide unit and sucrose octasulfate has 8 sulfate groups per disaccharide unit. This increased binding capacity for added cations means
- 20 that: a) more positively charged polyatomic complexes can be loaded onto the acidic saccharide matrix; and b) any stabilizing polyamine (polycation) preferably a polyamine such as protamine (protamine sulfate) which is typically added in minor quantity will bind the acidic saccharide
- 25 with higher avidity, as well as affinity. This is due to the higher number of anionically charged groups present on each saccharide unit, the closer proximity of one charged group to another. This gives rise to a greater possibility for multivalent binding between the
- 30 oppositely charged species of polyatomic complex and carrier. In specific examples, the  $\text{Cr}_4\text{S}^{+2}$  basic polyatomic species binds with higher avidity to negatively charged carrier acidic species. For example, such binding may be expected to involve chemical
- 35 cooperativity due to the close proximity (more densely spaced) of negative charges on pentosan polysulfate and

especially on sucrose octasulfate (both relative to heparin). Hence, the more densely charged the acidic groups, the greater the cross-linking of carrier with both the polyatomic complex and with added stabilizing (coacervating) polyamine (polycation) substances which include but are not limited to protamine (protamine sulfate) and hexadimethrine.

It will be evident to those skilled in the art, that any biologically compatible charged, divalent, multivalent or polyvalent substances which comprise acidic saccharides substances, polybasic substances such as polyamines, and respectively, their reciprocally charged groups or coacervating carrier substances can be used, respectively to prepare and further chemically stabilize those paired-ion (or coacervation stabilized) formulations. Specifically,  $Cr_4S$  polyatomic complex can be bound to negatively charged carrier substances, such as heparins, pentosan polysulfates and sulfated or carboxylated saccharides, including but not limited to sucrose octasulfate, sucrose pentasulfate, sucrose tetraacetate, glucose diphosphate, and other hexose, pentose and short-chain carboxylated, phosphated, and sulfated saccharides. Additional naturally occurring acidic saccharides include but are not limited to: dermatan sulfate, chondroitin sulfate, dextran sulfate, and hyaluronic acid, epithelial and other mucopolysaccharides, acidic saccharides and the like.

Conversely, it will be evident to those skilled in the art, that a positively charged (basic) carrier, including but not limited to protamine (protamine sulfate) and hexadimethrine can be used in conjunction with a negatively charged (or acidic) diagnostic actives (or drug actives). This and the carrier features which allows increased diagnostic active (or drug active)

loading capacity (see paragraph above), provide a basis for novelty and scope of the present invention. It will be evident to those skilled in the art that other classes of synthetic and naturally occurring polyamines can be  
5 used, including but not limited to polyamino acids, polylysine, polyarginine, histone proteins, heparin binding proteins, peptide fragments of the preceding substances, substances which contain more than one basic amino group, and the like.

10

A further novel feature described herein, is that closely spaced acidic groups, including but not limited to carboxylated and sulfated saccharides, and including but not limited to polyamines such as protamine and  
15 hexadimethrine, have the property of associating with and binding to determinants of vertebrate endothelium and epithelium. This property is envisioned as being further enhanced by formulations which embody a multivalent binding surface, including but not limited to, those  
20 formulations formed by assemblage of the paired-ion compositions, coacervates (multiply paired-ion compositions) and molecular complexes described above into nanoparticles or microparticles. To those skilled in the art, it will be evident that such endothelial  
25 association or binding to endothelium determinants, or to epithelial determinants, by the carrier substance or polyatomic complex-carrier combination, confers the additional potentially advantageous property of enhancing the internal physiologic or pathologic images related to  
30 either normal endothelium or diseased endothelium, or diseased tissues underlying the endothelium, provided that such disease renders its adjacent endothelium altered in some chemical or physical fashion. Hence, an additional broad property of endothelial association and  
35 binding, and in certain cases, transmigration, is

conferred by the simple and complex (coacervation) carriers envisioned herein.

The molecular weight ranges of carrier substances can be virtually any size below about 500,000 to 1,000,000 Daltons, above which size most macromolecules are no longer completely water soluble. For use in pharmaceutical formulations involving existing approved and new, synthetic and natural substances, the most typical, but not limiting, upper range of molecular weights of mutually binding (coacervating) substances is between about 22,000 Daltons (the upper MW range of standard heparins) and about 70,000 Daltons (the upper MW range of standard dextrans and dextran sulfates). This includes heparins of porcine source (typically ranging in size up to 22,000 Daltons), which, by virtue of the property of their longer chain length, may have increased propensity to self-associate even though each disaccharide unit carries a lower negative charge density relative to beef lung heparin, pentosan polysulfate or sucrose octasulfate. Because of granule characteristics, branched-chain configuration and exposure of potentially binding functional groups during processing, native and derivatized starches, including but not limited to carboxymethyl starch and hydroxyethyl starch, are also useful in forming coacervates and can also act as diagnostic ligand binding carriers and coacervate-forming substances, as can acidic and basic carbohydrates obtained from marine sources, including protans, carageenans, and the like.

The present Examples support mixtures of negatively and positively charged mixed carrier substances which contain as little as about 0.01% by weight of the minor component. The preferred weight ratio of protamine (protamine sulfate) to sucrose octasulfate is less than

about 2.0%, with the most preferred weight ratio being between about 0.12% and 0.20% (w/w of protamine). Weight ratios of less than about 0.5% protamine to pentosan polysulfate and protamine to heparin are preferred, with  
5 the most preferred ratios being in the range of about 0.1% to 0.2%, however, weight ratios outside this range are also envisioned as being preferred under special circumstances.

10 To those skilled in the art, the property of strong coacervation, paired-ion binding, multiply paired-ion binding and/or salt-forming properties of divalent or greater charged substances, when mixed together in  
15 combination, allows for an extensive combination of diagnostic and pharmaceutical actives and carrier substances to form chemically and/or physically stabilized formulations. It is envisioned that essentially any major-minor weight mixture of charge-complementary carrier substances which forms a stable  
20 complex or coacervate, will be useful in preparing the diagnostic formulations described in this application. Furthermore, such charge interaction is importantly envisioned as chemically or physically stabilizing the otherwise water-soluble separate components so that they  
25 remain strongly associated with one another in aqueous solvents and under physiological conditions, even in the absence of covalent chemical cross-linking or organic solvents.

30 A further novel feature described herein is the further chemical stabilization by opposite charges of the separate carrier components mixed together to form: molecular complexes, nanoparticles and microparticles. According to the most recent definitions, particles of  
35 nanometer dimensions, nanoparticles, are now further clarified as comprising particles of about 1.0 to 1,000

nanometers in diameter, and microparticles are now further clarified as comprising particles of greater than about 1.0 micrometer in diameter. For most in vivo parenteral diagnostic or pharmaceutical uses, the typical upper range of useful size for microparticles is about 15 micrometers.

Whereas it is envisioned that certain of the polyatomic complex carrier compositions herein will have preferred properties when formulated at less than about 5 percent (w/w) cross-linked or microaggregated species, it will be evident to those skilled in the art, that other physical formulations of the present invention, which are derived from the above, may take the form of nanoparticles or microparticles and still have significant advantages for selected in vivo diagnostic and pharmaceutical applications.

A further novel feature described herein is the additional stabilization, under both aqueous and dry-powder conditions, which is conferred by additionally drying the preceding formulations.

For pharmaceutical purposes, such drying is typically carried out by means of one or more of the following: spray drying, lyophilization (freeze drying), granulation drying, microwave drying, and the like. In drying, typically additional stability is conferred by conducting the drying process in the presence of amino acids and/or one or more of a sugar such as sucrose, mannitol, lactose, trehalose, and the like. In the present application, the weight percentages of such added sugars to polyatomic complex-carrier substances has been tested to range from zero (no added sugars) to about 1500% weight percent (in specific examples, lactose). However, to those skilled in the art, it is evident that

a wide range of concentrations of many biologically compatible polyols, sugars and/or amino acids which adds to the initial formulation stability, dry stability or stability following resuspension-redissolution, are  
5 useful for the present purpose.

A further novel feature of the formulations disclosed herein, is that combination of the polyatomic complex may, and typically does, confer increased dry-  
10 powder and resuspension stability upon the polyatomic complex, such that the combined or associated form of the complex has improved stability over time and/or during distribution in the bloodstream to sites of *in vivo* medical imaging. Hence, to those skilled in the art,  
15 particular ones of the present formulations which are most likely to be preferred are those for which the potency (R1 relaxivity -- see Examples and Table 5) are unchanged or increased in relation to the native polyatomic complex. In specific examples, this potency  
20 pertains to  $\text{Cr}_3\text{S}(\text{O}_2\text{CCH}_3)_6(\text{H}_2\text{O})_4^{+2}$ .

As an additional distinguishing feature of the present invention and its preferred embodiments, the mutual magnetic coupling of plural atoms with magnetic  
25 moments within (and only within) the polyatomic complex, together with the property of the bridging molecular species being free of intermolecular bonds, completely distinguishes the present, strongly paramagnetic (T1-relaxation) agents from externally superparamagnetic (T1-relaxation) agents from externally superparamagnetic  
30 agents, where the latter are defined by their magnetic hysteresis curves as having bulk (intermolecular or supramolecular) superparamagnetic properties (and being principally T2-relaxation agents). The latter include materials such as ferromagnetite,  $\text{Fe}_3\text{O}_4$ ,  $\text{Fe}_2\text{O}_3$ , and the  
35 like.

To those skilled in the art, it will be apparent that the novel features herein may be practiced optimally by combining the preceding formulations with standard pharmaceutical additives, including colorings, osmotically active salts and sugars, buffering substances including but not limited to monobasic and dibasic sodium and potassium phosphates, sodium hydroxide, hydrochloric acid, phosphoric acid, glycerol, polyethylene glycol, other polyols, amino acids and the like.

10

The following examples are presented to illustrate preferred embodiments of the present invention and their use in MR imaging or hysteresis heating. These examples are purely illustrative, and do not in any way delimit the full scope of the present invention.

15

#### EXAMPLE 1

##### PREPARATION OF DTPA-DEXTRANS

The cyclic dianhydride of DTPA (diethylenetriamine pentaacetic acid), as prepared by the method of Eckelman et al. (J. Pharm. Sci. V 64, pp 704-706 (1975), was obtained in a highly pure form from Calbiochem-Behring Corp. The completely soluble DTPA derivative of dextran was prepared by adding 7.0 g of the cyclic DTPA dianhydride stepwise to 1.72 g of Dextran T70 (average MW 70,000 Daltons,  $M_n$  46,000, Pharmacia Chemicals) in a reaction solvent comprising HEPES buffer 115 mg/100 cc distilled water, pH 7.0 to 8.0. The reaction was carried out with vigorous stirring at ambient temperatures for 1 hr with readjustment to pH 7.0 using NaOH, after the addition of each aliquot of DTPA dianhydride.

30

The dextran-DTPA product was separated from unconjugated DTPA by dialysis against 200 volumes of 0.15 N NaCl and then 50 volumes of distilled water at pH 6.5. This and the following step are major improvements over

35



the derivatization method previously filed by the present inventor. (See U.S. Patent Applications 799,757 and 086,692, and PCT Application PCT/US86/02479, which are hereby incorporated by reference.) Upon completion of  
5 dialysis, the conjugate was brought again to 115 mg/100cc in HEPES buffer, and reacted a second time with an identical quantity of DTPA dianhydride as described above. After this, the dialysis was repeated as described above.

10

As assessed by molecular filtration, 98% of the dextran-DTPA product had a molecular weight of less than 100,000 Daltons. The dilute solution of DTPA-dextran was either: a) concentrated to between 10% and 25% (w/v) by  
15 forced filtered-air evaporation at room temperature, or b) lyophilized to dryness for prolonged shelf storage. Concentrated salts and buffers were added as needed, to render the final preparations physiologically acceptable for injection. The pH was maintained between 6.5 and  
20 7.0. As assayed by complexometric titration, one ligand of DTPA was conjugated for every 7 sugar residues, for a total of 55.5 DTPAs per 389 glucose units in each average molecule.

25

Two other soluble DTPA-dextran derivatives were synthesized from dextrans of starting molecular weights = 10,000 Daltons (Dextran T10, Pharmacia Chemicals) and 40,000 Daltons (Dextran T40, Pharmacia Chemicals). All of the preceding dextrans were soluble and free of  
30 microaggregates, as assessed by filtration through serial molecular sieve filters (Amicon Corporation).

## EXAMPLE 2

### PREPARATION OF DTPA-HYDROXYETHYL STARCH

35

Low-molecular-weight hydroxyethyl starch is obtained in a highly pure and soluble form from American Critical

Care/DuPont, reacted with the cyclic dianhydride of DTPA, and the polymeric derivative separated, concentrated and titrated as described in EXAMPLE 1.

5 EXAMPLE 3

PREPARATION OF SUCCINYLATED-DEXTRANS

Succinyl anhydride is obtained in a highly pure form from Aldrich Chemicals, and reacted with dextrans of 40,000 MW and 70,000 MW, and the polymeric derivative  
10 separated, concentrated and titrated as described in EXAMPLE 1.

EXAMPLE 4

PREPARATION OF THE PAIRED-ION

15 METAL CO-ORDINATE-POLYMER

COMPLEX CISPLATIN-DTPA-DEXTRAN (70,000 MW)

Lyophilized DTPA-dextran (70,000 MW), prepared as in Example 1, was dissolved in 1.4 cc of sterile water, heated for 30 seconds by swirling in a boiling water  
20 bath, added (hot) to 70 mg of Platinol™ powder (containing 3.33 mg of cisplatin,  $\text{Pt}(\text{NH}_3)_2\text{-Cl}_2$  with the remaining weight comprising excipients; Bristol Laboratories), the solution vortexed vigorously for 30 seconds to dissolve the Platinol™ + excipients, and the  
25 resulting solution cooled to room temperature and checked for complete solubility at 2.4 mg/ml. Formation of a stable paired-ion complex between the platinum coordinate and the carboxyl groups bound to dextran, was established by three tests: a) continued solubility of cisplatin at a  
30 concentration greater than its native solubility limit of 1.5 mg/cc; b) reduction in the complexation of exogenously added calcium ions by the DTPA groups of DTPA-dextran (assessed using an Orion Instruments ionized calcium analyzer); and c) elimination of tetany following  
35 intravenous injection of the resulting mixture into CBA/J mice (Jackson Laboratories) at a dose of 10 mg/25 gm body

-69-

weight. This absence of tetany contrasts with the occurrence of tetany and death in mice which were injected with an equivalent dose of DTPA-dextran alone - and importantly, in the absence of balancing quantities of calcium ion, which render the resulting Ca-DTPA-dextran entirely nontoxic *in vivo*. Hence, by *in vitro* and *in vivo* criteria, cisplatin (as Platinol™) undergoes complexation to DTPA-dextran at a sufficient binding stability to compete with an added, divalent metal cation ( $\text{Ca}^{+2}$ ). The exact coordination state and chemical structure of the resulting cisplatin-DTPA-dextran complex has not been further elucidated.

#### EXAMPLE 5

##### 15 PREPARATION OF THE PAIRED-ION METAL COORDINATE-POLYMER COMPLEX, CISPLATIN-HEPARIN (22,000-26,000 MW)

Beef-lung heparin (Upjohn Company) 6,000-10,000MW was added dry at 14 mg to 280 mg of dry Platinol™ powder (Bristol Laboratories) containing 14 mg of cisplatin, the mixture dissolved in 14 cc of sterile water and vortexed for 1 minute to completely dissolve all components. Formation of a stable paired-ion complex between the platinum coordinate and the sulfate groups covalently bound to heparin, was established by two tests: a) continued solubility of cisplatin at a higher concentration (2.25 mg/cc) than its native solubility limit of 1.5 mg/cc; and b) alteration of cisplatin biodistribution following intravenous and intraarterial injection in animals (see the Examples below).

30

EXAMPLE 6PREPARATION OF HEAT-STABILIZED, HYDROXYETHYL STARCH-MATRIX MICROPARTICLES WHICH ENCAPSULATE CISPLATIN AND HAVE A HEPARIN SURFACE COATING

5        Hydroxyethyl starch 605 mg (Sigma Chemicals) was  
suspended in 5.5 cc of sterile water and heated for 3  
minutes in a boiling water bath to bring it into a stable  
(translucent) emulsion, and 5 cc of this was added to  
1000 mg of Platinol™ (Bristol Laboratories) containing 50  
10 mg of cisplatin. This was emulsified for 30 seconds in  
70 cc of heated (100°C) cottonseed oil (Sargent Welch)  
using a Brinkmann Instruments ultrasonic homogenizer, and  
the oil cooled in a room-temperature water bath, with  
continued homogenization for 2 more minutes, until the  
15 emulsion itself reached room temperature. This was  
extracted 4 times with acetone (Fisher Chemicals)  
containing 0.5% Tween 80 (Sigma Chemicals), and was  
harvested by centrifugation and air dried. The resulting  
microparticle diameters ranged from 0.1-1.0 micron.

20        A heparin coating was applied to the particle  
surfaces by adding 2 cc of a water solution containing 50  
mg of beef-lung heparin (Upjohn Company) 6,000-10,000 MW,  
adding the particle suspension plus heparin to 70 cc of  
25 heated (100°C) cottonseed oil and repeating the  
emulsification and extraction steps described in the  
preceding paragraph. The resulting particles ranged from  
0.1 to 0.8 micron in diameter. The presence of a heparin  
surface coating was verified by suspending the particles  
30 in normal saline and adding protamine (Sigma Chemicals),  
a multivalent heparin-binding agent. This produced  
aggregation and agglutination of the heparin-coated (but  
not uncoated) particles.

EXAMPLE 7IN VIVO TESTING FOR SELECTIVE LUNG LOCALIZATION OF THE  
PRECEDING PREPARATIONS FOLLOWING INTRAVENOUS  
ADMINISTRATION

5 CBA/J mice (Jackson Laboratories) were injected intravenously via the tail vein with a) small microparticles containing a heparin surface (as in Example 6) and b) soluble heparin-cisplatin complex-es (prepared as in Example 5). At 5 to 15 minutes  
10 postinjection, the animals were sacrificed, their lungs removed and fixed with intratracheal buffered formalin, tissue sections cut at 8 microns thickness, and the sections stained using a newly devised method for platinum which comprises a microwave-augmented iron-type  
15 stain (60°C x 2 minute x 3 cycles) using a 1:1 mixture of 2% ferriferrocyanide reagent and 4% HCl. By this method, lung uptake of both the small microspheres (see Figure 4) and soluble paired-ion complex of cisplatin-heparin was documented at the 5-minute postinjection interval. Rapid  
20 uptake occurred in both extracellular and intra-cellular compartments, and additional histochemical positivity of bronchial respiratory epithelium and paratracheal lymph nodes was observed at 10-15 minutes. No significant staining was observed following intravenous injection of  
25 a standard formulation of Platinol (Bristol Laboratories). To those skilled in the art, additional evidence for selective lung localization, was obtained by injecting intravenously, analogous (subembolizing) heparin-coated small microparticles containing encapsul-  
30 ated amphotericin B, into identical mice, homogenizing the lungs, and documenting an 8-fold increment in drug levels over native amphotericin B (deoxycholate formulation, Fungizone; Squibb) recovered at 1 to 3 hours postinjection. (See U.S. Patent Application No.  
35 07/033,432, and PCT application PCT/US88/01096, which are hereby incorporated by reference.) Hence, the preceding

histologic stains correlated with an increment of nearly 1 order of magnitude in selective pulmonary carrier and drug localization.

5 EXAMPLE 8

IN VIVO TESTING FOR SELECTIVE TUMOR LOCALIZATION OF THE PRECEDING PREPARATIONS FOLLOWING INTRAARTERIAL ADMINISTRATION

Additional documentation for maintenance of the cis-platin-carrier paired-ion complex *in vivo* was obtained as follows. Rabbits bearing VX2 carcinomas of the right hind limb were catheterized under fluoroscopic control, and three of the preceding Platinol formulations, as well as standard Platinol, were injected at a constant dose of 15 mg (of cisplatin) per rabbit by selective arterial perfusion over 15 minutes, into the tumor-bearing limb. Animals were sacrificed at 15 minutes, and the tumors and organs were homogenized extracted and analyzed by atomic absorption for tissue platinum concentrations, as shown in Table 1.

Table 1  
Platinum content (ng/mg of tissue, wet weight) -  
Ipsilateral

<u>Agent</u>	<u>Blood</u>	<u>Tumor</u>	<u>Muscle</u>	<u>Liver</u>	<u>Kidney</u>
Heparin-cisplatin:	2.71	12.24	0.18	5.74	5.88
DTPA-dextran cisplatin:	2.14	10.81	0.29	3.47	2.92
Heparin-coated hydroxyethyl starch small microspheres of cisplatin:					
	2.43	14.07	0.20	4.64	5.91
Standard:	2.36	8.40	1.09	6.09	4.09

Additionally, histochemical platinum stains were performed on the tumor tissues, as described in Example 7. These stains revealed intracellular tumor-cell (but

-73-

not normal-cell) platinum in all of the groups in Table 1 except group 4. Moreover, the intracellular staining of tumor cells in groups 1-3 was significantly more intense than the background staining of hemoglobin iron in red blood cells. Since hemoglobin iron is present at a mean corpuscular hemoglobin concentration (MCHC) of about 0.2 molar, these results suggest strongly that tumor cell platinum reaches very high concentrations relative to those achieved with standard Platinol™. To individuals skilled in the art, this also indicates that selective tumor-cell augmentation of hysteresis heating may be achieved by first administering one or more of the carrier formulations described above, but in stable complexation with a strong superparamagnetic substance which is otherwise too small and uncontrolled to undergo this degree of selective localization.

#### EXAMPLE 9

#### PROLONGED ENHANCEMENT OF HUMAN TUMOR (MELANOMA) XENOGRAPHS IN NUDE MICE, BY PARAMAGNETIC CHELATE ASSOCIATED COVALENTLY WITH DEXTRAN 70

The strong paramagnetic metal ion, gadolinium ( $Gd^{+3}$ ), was chelated stoichiometrically to DTPA-Dextran 70 polymer, whose preparation is described in Example 1, paragraph 2 (improved formulation). Swiss nude mice were inoculated with BRO-strain human malignant melanomas, and these were allowed to grow to a 1-1.5 cm diameter. Moderately T1-weighted MR imaging ( $TR = 500$  msec,  $TE = 40$  msec) was performed in a standard Diasonics™ medical imager and a 30-cm RF head coil, before and after intravenous injection of equivalent doses of Gd-DTPA-Dextran-70 (0.03 mmol Gd/kg) or Gd-DTPA (0.1 mmol/kg) contrast agent. Gd-DTPA-dextran and Gd-DTPA began to optimally enhance the tumors at comparably short postcontrast intervals of 10 minutes, however, by one hour Gd-DTPA had completely faded, whereas Gd-DTPA-

dextran continued to enhance these tumors intensely for longer than 2.5 hours (the cutoff time on imaging experiments). To those skilled in the art, it will be recognized from these results that chelated Gd and small metal coordinates in general, will benefit greatly in terms of potency (by at least half an order of magnitude), tumor selectivity and tumor retention, from covalent conjugation or strong paired-ion association with dextran or including analogous carbohydrate carrier molecules. It will also be recognized that much lower doses of the stronger superparamagnetics (including doses below about 0.01 mmol/kg, versus 0.1 mmol/kg for Gd-DTPA) can be used to obtain effective MRI contrast enhancement; and that much faster, more heavily T1-weighted pulse sequences can be implemented in the presence of intramolecularly superparamagnetic polyatomic complex-carrier polymer conjugates or paired-ion complexes.

In the preceding study (see PCT Patent Application PCT/US88/01096),  $^{153}\text{Gd}$ -DTPA-dextran was documented to clear from the blood with a  $t_{1/2}$  of about 50 minutes (versus 20 minutes for Gd-DTPA). Total body clearance was almost complete (greater than 96%) by 24 hours. Hence, to those skilled in the art, it will be recognized that strong association of a metal coordinate with this polymeric carbohydrate of predominantly less than 50,000 MW, and particularly with a polymeric carbohydrate with substantially lower molecular weight of less than 10,000 Daltons, allows rapid and complete blood and body clearance by predominantly renal pathways.

#### EXAMPLE 10

PREPARATION OF PAIRED-ION MOLECULAR COMPLEXES OF  
 $\text{Cr}_2\text{S}(\text{O}_2\text{CCH}_2)_2(\text{H}_2\text{O})^{+2}$  WITH NEGATIVELY CHARGED

POLYMERIC CARBOHYDRATE CARRIERS



-75-

The following negatively charged polymeric carriers are obtained for individual addition and ion pairing to the ferromagnetically coupled polyatomic cation complex  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  (Exxon Corporation) (see the Bino et al. article cited above): heparin (6,000-10,000 Daltons, Upjohn Company); DTPA-dextran (40,000 and 70,000 MW parent carbohydrates, derivatized as in Example 1); DTPA-hydroxyethyl starch (50,000 MW parent carbohydrate, prepared as in Example 2); and succinylated-dextran (40,000 and 60,000 MW parent carbohydrates, derivatized as in Example 3). Each polymer is added as a concentrated aqueous solution, at stoichiometric charge equivalency, or at 50% or 25% of charge equivalency, to a concentrated aqueous solution of  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  as the chloride salt. Ion pairing is achieved by direct mixing and heating for 1-5 minutes to 100°C at pH 7, with vigorous stirring. To subfractions of the 25% and 50% mixtures (above) is added a concentrated aqueous solution of N-methylglucamine at quantities sufficient to achieve electrical neutrality. The stability of ion pairing is tested by performing equilibrium dialysis against 200 volumes of 0.15 N NaCl and assaying the retained (polymeric) materials for T1 relaxivity (IBM PC20 NMR Spectral Analyzer). Those skilled in the art will recognize from the results of cisplatin complexation to heparin and DTPA-dextran (documented in Examples 4 and 5) that the even more positively charged  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  counterion of the present example results in an even stronger ion pairing to heparin and DTPA-dextran than the satisfactory (*in vitro* and *in vivo*) pairing achieved for the cisplatin metal-amine coordinate documented in Examples 4 and 5.

EXAMPLE 11

PREPARATION OF  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$   
BOUND TO HEPARIN

Beef-lung heparin Fraction A (6,000-10,000MW) is obtained (Hepar Corp.) for individual binding to the polyatomic chromium complex,  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4\cdot\text{Cl}_2$  (Chiron, Trondheim, Norway). Since the chromium complex is relatively unstable to prolonged heating in water, this reaction is carried out over a short interval in absolute ethanol (USP) (in which the chromium complex shows markedly increased stability to heating) and under increased pressure, using a microwave digestion bomb (Berghof America). The heparin is sonicated into the ethanol as a very fine solid dispersion. The chromium complex is added under concentrated conditions of 250 mg/ml, in order to take advantage of its autoprotection at higher concentrations against degradation in hot solvents. This mixed dispersion/solution is placed in the digestion bomb, the bomb is sealed, the sealed bomb is placed into a preheated, 135C glycerol bath for 5 minutes, the reaction is carried out with very rapid magnetic stirring, and then the bomb is transferred into water at room temperature in order to achieve rapid cooling to ambient temperature. The resulting product remains as a very fine dispersion whose size varies depending on the size of the heparin at input. This size ranges from about 100 nanometers to about 10 micrometers in mean diameter. The resulting products respectively, are centrifuged at 22,000 x g for 10 to 15 minutes, and binding of the chromium complex together with retention of at least partial integrity (greater than about 50%) of the chromium complex are verified by UV measurements at the characteristic maxima of 600 nanometers.

#### EXAMPLE 12

PREPARATION OF  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4\cdot\text{Cl}_2$

BOUND TO DIETHYLENETRIAMINEPENTAACETATE HEPARIN

AND TO DIETHYLENETRIAMINEPENTAACETATE DEXTRAN

Beef lung heparin Fraction A (6,000-10,000 MW) is obtained (Hepar Corp.) for individual binding to polyatomic chromium complex,  $[\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4]\cdot\text{Cl}_2$  (Chiron, Trondheim, Norway), and is dissolved in distilled water at 25 mg/ml.

Diethylenetriaminepentaacetate (DTPA) heparin and DTPA dextran are prepared as in Example 1, and are added respectively, to the chromium complex in aqueous solution, such as to give a molar ratio of 0.4 moles of DTPA carrier to 1.0 moles of chromium complex. The respective reactions are carried out by placing the reaction tubes into a beaker of boiling water for preferably 60 to 90 seconds, and then immediately quenching the reaction by transferring these reaction tubes into a beaker of room temperature water. Evidence for transchelation binding of the chromium complex by carrier-bound DTPA comprises: a) molecular filtration retention of the characteristic blue color above a 3,000 Dalton molecular cutoff filter; and b) a downward shift in the UV absorbance maximum of the resulting product from the native 600 nm peak of native chromium complex, to about 560 nm. A slight to moderate decrease in absorbance intensity is observed, suggesting that some degradation of the chromium complex accompanies this reaction. However, such degradation is not to that of comparably heated chromium complex alone, which retains an absorbance maximum of 600 nm. Additionally, such degradation is not to that of chromium complex alone, comparably heated, or of chromium complex plus a two-fold molar excess of DTPA comparably heated, both of which have characteristically distinct UV absorbance maxima and absorbance peak height ratios for their characteristic two absorbance maxima at about 500-600 nm and about 390-450 nm. Hence, there is strong evidence for formation of stabilized polymeric chromium complex-heparin and chromium complex-dextran 70 agents, wherein binding is by

partial transchelation of the chromium complex to the DTPA groups of the carriers. Further evidence involves preservation of at least about 60-80% of the native R1 relaxivity of the free chromium complex.

- 5        Figure 1 shows a perspective view of the  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  cation (excluding hydrogen atoms). The atoms are shown with ellipsoids, to indicate approximate thermal vibration ranges at room temperature.

10    EXAMPLE 13

ADVANTAGES TO NMR IMAGING OF POLYMERIC FORMULA-  
TIONS OF CARRIER FERROMAGNETICALLY COUPLED POLYATOMIC  
COMPLEXES

- Selected polymeric formulations of
- 15     $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ , prepared as described in the preceding Examples, are injected intravenously to obtain systemic lesional uptake (frequently practiced in radiology), or intraarterially (less frequently practiced in radiology), to obtain highly selective uptake in
- 20    regional tumors, especially of the liver, pelvis, brain and limbs. Those skilled in the art will recognize that the more potent, more selective, less toxic (including especially chromium nontoxicity) polymeric formulations of superparamagnetics, including  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  and
- 25    analogous cluster compounds incorporating ions such as  $\text{Gd}^{+3}$  or  $\text{Fe}^{+3}$ , advantageously allow the dose of paramagnetic, including chromium, to be reduced to less than about 0.005 to 0.01 mmol/kg of body weight. (See Ranney, Contrast Agents in Magnetic Resonance Imaging, in
- 30    Excerpta Medica at page 81 (1986), which is hereby incorporated by reference). It will also be recognized that MRI procedures involving fast imaging (see Bluem et al., 157 Radiology 335 (1985), which is hereby incorporated by reference) and cardiovascular MRI
- 35    (including MRI "angiography") (see Nagler et al., 157 Radiology 313 (1985), which is hereby incorporated by

reference) will benefit greatly, in terms of shortened image acquisition time and improved quality, from a non-toxic contrast polymer of super-paramagnetic potency. Additionally, the polymeric polyatomic-complex agents described in the preceding Examples are useful for co-labelling either therapeutic drug carriers (polymeric or nanoparticulate) or the therapeutic agents themselves, whose tumor (or other lesional) localization needs to be monitored and whose rate of release from the carrier (bioavailability) needs to be assessed noninvasively *in vivo*, potentially in multiple lesions at different depths within a body region. In this context, polymeric polyatomic-complex agents are useful and of improved utility due to increased potency and selectivity, and reduced toxicity. Furthermore, increased potency allows drug release to be monitored over longer postinjection intervals within target tissues, organs, tumors and infections. These improvements are based on the present application, and also, in part, on applicant's earlier-reported work on the use of partially analogous, but less potent, Gd-DTPA-dextran-labelled drug carriers leading to localization in tissues and enhanced MRI detection. (See the Ranney and Huffaker article at 507 Proc. NY Acad. Sci. 104 (1987), which is hereby incorporated by reference.)

#### EXAMPLE 14

#### ADVANTAGES TO HYSTERESIS HEATING OF POLYMERIC FORMULATIONS OF FERROMAGNETICALLY COUPLED POLYATOMIC COMPLEXES

Microinhomogeneities of tissue heating represent a major problem in hyperthermia treatment of tumors. This results in considerable part, from the selective survival of tumor cells lying adjacent to microvessels - in which heat loss is accentuated by blood flow. A partially effective approach to this problem has been to inject

small ferromagnetic particles of  $\text{Fe}_2\text{O}_3$  directly into the tumor masses, and then apply magnetic hysteresis heating at frequencies of 10-100 kHz to the entire local region. (See Borelli et al., 29 Phys. Med. Biol. 487 (1984).

5 Effective superheating and tumor regression in mice occurs if the injected magnetic material is present 1) in sufficient quantity and 2) at a sufficient macrodomain size for efficient hysteresis augmentation to occur (including 0.5-2 micron particle diameters). Because the

10 carrier/polyatomic complex agents and ion-cluster agents described in the preceding Examples have the properties of markedly improved selectivity and dose of tumor localization, retention in the viable (perfused) subregions of tumor, and improved tumor-cell uptake, it

15 will be understood by those skilled in the art that these carrier/polyatomic complex and ion-cluster agents can be of significant benefit in augmenting the homogeneity, magnitude and tumor-cell selectivity of hysteresis heating induced by oscillating magnetic fields, provided

20 that the associated superparamagnetic agents (which may be associated by conjugation, ion-pairing, or encapsulation) become concentrated as adequately sized macrodomains (of at least about 0.5 micron) in the target sites or cells. Histologic staining for cisplatin (per

25 Example 7) of the VX2 rabbit carcinomas which were perfused with heparin-coated cisplatin-hydroxyethyl starch small microspheres (of 0.1-0.8 micron diameters), documented that many of the tumor cells in the target region addressed by selective arterial perfusion, stained

30 intracellularly in a punctate pattern, wherein the diameters of punctate staining positivity ranged from 0.1 to 0.8 micron (= the diameters of the original particles). Importantly, high (including about 0.2 molar) intracellular concentrations of cisplatin were

35 achieved in the VX2 carcinoma cells *in vivo* (see Example 8). The combination of these high levels plus

intracellular aggregation were achieved by administering the cisplatin formulated as a heparin-coated microparticle. This documents the type and extent of intracellular accumulation and aggregation of super-  
5 paramagnetics which can be achieved and are potentially useful in locally amplifying exogenous hysteresis heating. Notably, aggregated staining was absent following intraarterial perfusion of standard (soluble, low-molecular-weight) cisplatin. Hence, in the present  
10 example, heparin-coated  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  hydroxyethyl starch small microspheres of 0.1 to 0.8 micron and intermediate-sized microspheres of 0.8-3.0 micron diameters, when administered intraarterially, are useful as amplifiers of hysteresis heating, and therefore as  
15 inducers of augmented cell death *in vivo*.

#### EXAMPLE 15

##### NOVEL GLYCINE-SUBSTITUTED POLYATOMIC Cr COMPLEX

The substrate,  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4$ , was prepared as  
20 described by A. Bino et al. (*Science*, Vol. 241, pp. 1479-1482, Sept. 16, 1988). This was added at 0.5 gm to 50 cc of acetic anhydride, followed by the addition of 0.1 gm of glycine. The mixture was refluxed at 142 degrees C for 4 hours, resulting in a yellow-green solution which  
25 was poured while still hot into cold water. Evaporation of the solvent gave yellow crystals, which, when recrystallized, gave an infrared (IR) spectrum indicative of glycine substitution for some of the acetate bridging groups, and retention of Cr-O and Cr-S bonds (Fig. 5C, with Figs. 5D and 5E representing the appropriate  
30 controls). Specifically, the IR bands around  $1600\text{ cm}^{-1}$  (Fig. 5C) are characteristic of ionized bidentate glycine, and the bands between  $300$  and  $400\text{ cm}^{-1}$  are also consistent with this. The additional bands between  $300$   
35 and  $400\text{ cm}^{-1}$  are also consistent with the presence of Cr-O and Cr-S bonds. Figs. 5A and 5B show the degradation

-82-

controls wherein the  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4$  degrades to ionic chromium (by IR spectral criteria) when heated with glycine in an aqueous (protonating) solvent, water, for 4 hours at 92 degrees C. These results provide strong evidence for the formation of a novel polyatomic chromium-atom cluster compound of the general formula,  $\text{Cr}_n\text{S}(\text{O}_2\text{CCH}_3)_x(\text{Gly})_y(\text{H}_2\text{O})_z$ , wherein n is the number of Cr atoms greater than 1, x is the number of acetates between about 3 and 7, y is the number of glycines between about 1 and 5, and z is the number of loosely bound waters. The inclusion of bidentate glycine as a novel bridging ligand in the polyatomic chromium cluster provides a new reactive (charged) group for binding to carriers and renders the entire compound novel.

#### EXAMPLE 16

##### NOVEL POLYATOMIC GADOLINIUM CLUSTER COMPOUND

Gadolinium chloride (3.7 gm, Alpha) and  $\text{Na}_2\text{S}$  (2.4 gm, Sigma) were mixed in 25 cc of a 1:1 mixture of glacial acetic acid and acetic anhydride and refluxed at 138 degrees C for 3 days. After removal of the dark red, gadolinium-negative filtrate, a lightly tan-colored precipitate was recovered which was qualitatively positive for gadolinium and which, after recrystallization, yielded a water-soluble water-stable compound whose IR spectrum was consistent with gadolinium acetate. Importantly, several additional bands were present in the carbonyl (ca.  $1500\text{ cm}^{-1}$ ) and C-O (ca.  $1000\text{ cm}^{-1}$ ) bond stretching wavelengths, which are strongly indicative of a polyatomic (polymeric) nature of this complex and, hence, indicative of a novel, gadolinium-containing, polyatomic complex for use with the disclosed carriers.

#### EXAMPLE 17

##### FORMATION OF STABLE CARRIER/ POLYATOMIC METAL ATOM COMPLEXES



Carboxymethyl dextran was obtained commercially and mixed with  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4$ . The resulting complex had a T1 relativity (R1) 3 times greater than that of the simple chromium complex, indicative of paired-ion binding between the chromium complex and the carrier. Analogous mixtures were performed of  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4$  and (a) dicarboxyethyl dextran; (b) heparin; and (c) dextran sulfate. The resulting paired-ion complexes were stable to dialysis in 0.15 molar (isotonic) to 0.5 molar saline.

10 EXAMPLE 18

STABILIZATION OF CHROMIUM COMPLEX AND COMPLEX-CARRIER ASSOCIATIONS. AS MICROPARTICLES AND NANOPARTICLES BY DRYING AND HEATING IN ORGANIC SOLVENTS

A. Chromium complex was obtained as in Example 11, dissolved in water at 25 mg/ml, Fraction A bovine lung heparin (6,000-8,000 MW) was added at a weight ratio of 4 parts heparin to 1 part chromium cluster, and this mixture was adjusted with NaOH to between about pH 5.5 and pH 7.0, and then magnetically stirred for 1 to 24 hours at 4C, with resultant formation (by interference light microscopy) of 1-4 um dark blue microparticles. This material was lyophilized for 24 hours in the presence and absence of mannitol plus dextrose in equimolar ratio, with the combined excipients equalling 300% by weight of the chromium complex-heparin microparticle weights. Upon resuspension of the lyophilized particles and examination by laser light scattering, only moderate transient physical stability of the particles was observed, with retention of nanoparticle and microsphere sizes for about 5 minutes.

Hence, these lyophilized materials were further resuspended respectively, in polyethylene glycol, glycerol, cottonseed oil, corn oil, acetone, acetone containing polyoxyethylene sorbitan mono-oleate at about 0.01% to 25% by weight to the lyophilized materials, and these materials were dispersed in the organic phase

either with low energy mixing or with high energy (pressure) dispersion methods, including rotor-stator dispersion and probe and bath sonification (= sonication). The materials were then heated, respectively at temperatures ranging from 110°C to about 140°C, for between 3 and 5 min. For acetone formulations, heating were carried out under increased pressure in a microwave digestion bomb (Berghof America) with magnetic stirring. Depending on the type and extent of energy input and heating, this resulted in formation of blue nanoparticles of about 45 to 85 nanometers in mean diameter (Nicom laser light scattering sizer), or blue microparticles of about 0.5 to 5.0 micrometers in mean diameter (Coulter Multisizer). Zeta potential measurements (Coulter Delsa 440) indicated that both the nanoparticles and microparticles had moderately negative surface charges of between about -40 and -50 millivolts. Ultraviolet absorbance measurements indicated that a majority of the characteristic 600 nm absorbance of chromium complex was retained for the particulate products obtained in all organic solvents, and that acetone-based solvents resulted in the highest preservation ratios to the native complex. The resulting particles were stable in distilled water and 55 dextrose in water, for intervals of between about 15 minutes and 48 hours after resuspension.

B. Chromium complex was obtained, dissolved, pH-adjusted, mixed with heparin and incubated with stirring as described in A. (above) but was not lyophilized. The complex-carrier adduct which formed was precipitated out of aqueous solution by stepwise addition of acetone to a final volume ratio of 2:1 (acetone:water). This material was resuspended by sonification in acetone containing polyoxyethylene sorbitan mono-oleate (at weight ratios as described in A., above) and the resuspended material was

heated to between 110°C and 140°C, under increased pressure in a microwave digestion bomb, as described in A., above). The resulting product was harvested by either air drying or lyophilization. Upon resuspension, the resulting nanospheres and microspheres retained their characteristic blue color and were stable to resuspension in water and 5% dextrose in water for up to 72 hours.

#### EXAMPLE 19

#### STABILIZATION OF CHROMIUM COMPLEX AND COMPLEX-CARRIER ASSOCIATIONS, AS MICROPARTICLES AND NANOPARTICLES BY COMBINATION WITH SECOND CARRIER (MATRIX) MATERIALS

Fraction A beef-lung heparin of between about 6,000-10,000 MW (Hepar Corp.) was dissolved at 100 mg/ml in distilled water, and this was added, respectively, to protamine sulfate (Sigma) or hexadimethrine (Sigma) predissolved in water at 15 mg/ml, such that the weight ratios varied from 2% to 98% heparin (relative to the second carrier material). This resulted in stable, characteristically blue nanoparticles and microparticles, which retained their physical particulate stability (without heating) in water and in 5% dextrose in water, for intervals of up to and greater than 24 hours, for nanoparticles and microparticles, respectively. Protamine induced stability is preferred and gives physical stabilization for greater than 24 hours. The optimal ratio of protamine is between about 2.5% and 15% (w/w). Both the microparticles and nanoparticles could be harvested by centrifugation at 22,000 x g for 10 minutes. The resulting nanoparticles so harvested, retain greater than about 60% of their starting magnetic potency, as evaluated by R1 relaxometry (IBM PC20 Relaxometer), and the microparticles retained greater than about 40% of the starting magnetic potency (same method). The differences in potency were consistent with greater availability of magnetic materials to diffusible

water when these materials are comprised as smaller nanoparticles than as larger microparticles.

EXAMPLE 20

TESTING FOR EXCLUSION OF CROSS-LINKED

5 AND MICROAGGREGATED SPECIES

IN THE SOLUBLE POLYMERIC

POLYATOMIC CHROMIUM COMPLEX-CARRIER FORMULATIONS

The soluble polymeric intermediates and final formulations prepared in previous Examples 11, 12, 18 and  
10 19 above, are tested for absence of cross-linking and microaggregation by ultrafiltration through a 10,000 MW cutoff filter (Amicon Corp.). In each case, less than about 5% of the total material of the soluble agents is retained above the filter.

15

EXAMPLE 21

TESTING FOR RELATIVE IN VITRO POTENCIES  
20 OF BOVINE AND PORCINE FRACTION A HEPARINS

Fraction A heparins of between about 6,000 and 10,000 MW are obtained from bovine lung and porcine mucosal sources respectively (Hepar Corp.), these are added at increasing standard concentrations to Azure A  
25 dye (Sigma), and the resultant decrease in the characteristic UV absorbances at 620 nm are determined. The absorbances decrease linearly with heparin concentration over a range of about 0.1 to 10 ug/ml (and per assay tube), however, the slope for bovine lung  
30 Fraction A heparin is approximately 15 percent steeper than that for porcine mucosal Fraction A heparin. This provides in vitro correlative evidence that porcine mucosal heparin is less potent than bovine lung heparin in an in vitro binding assay (Azure A) which correlates  
35 empirically with both (published) differences in sulfation ratios (bovine being greater) and in vivo lung endothelial binding and targeting (see below).

EXAMPLE 22TESTING FOR RELATIVE IN VIVO POTENCIES OF BOVINE AND  
PORCINE FRACTION A HEPARINS

5        Fraction A bovine lung and porcine mucosal heparins  
of between about 6,000-10,000 MW are obtained as in  
Example 21, and are used to prepare microparticulate  
dispersions of the therapeutic agent, amphotericin B, for  
10        which a rapid high performance liquid chromatographic  
ultraviolet detection assay exists. The microparticulate  
dispersions of both heparin formulations have a mean size  
of about 5 to 8 micrometers, as assessed by interference  
light microscopy. At 1 hr following intravenous  
15        injection into Balb/c mice and  
homogenization/solubilization of the lungs for  
determination of lung amphotericin B levels, there is a  
substantial difference in the lung localization of  
amphotericin B core material which appears to result from  
differences in binding of the different heparin surface  
20        materials, as follows:

	Microparticles prepared from	Lung amphotericin
	B	
	1. Bovine lung heparin	27.6 ug/gm
25	2. Porcine mucosal heparin	6.0 ug/gm

Although this result is obtained for core material  
other than the MRI enhancing agent material described in  
the current examples, to those skilled in the art, it  
30        indicates strongly that Fraction A heparin from bovine  
lung source is significantly more active for endothelial  
binding, transport and lung uptake than are the same  
Fraction obtained from porcine mucosal source, although  
it should be noted that porcine Fraction A heparin does  
35        indeed have endothelial binding/transport/uptake  
activity. This provides the initial in vivo basis for

the present new, novel formulations for endothelial binding, transport and tissue uptake based on low molecular weight (less than about 10,000 MW) Fraction A heparins, and in particular on such fraction a heparin  
5 obtained from bovine lung source.

EXAMPLE 23

IMAGING DATA

Preparation of  $(Cr_4S(O_4CCH_3)_6(H_2O)_6)$ -heparin carrier  
(hereinafter "Cr<sub>4</sub>S-heparin") for in vivo Magnetic

10 Resonance imaging

Two aliquots of the  $Cr_4S$  ion cluster (37.5 mg each) were dissolved in 1 ml (each) of 5% dextrose in water. To the first aliquot was added 187.5 mg of dextran sulfate carrier (8000 Daltons); and to the second aliquot  
15 was added 187.5 mg of beef-lung heparin carrier (ca. 6,000-10,000 MW Daltons). Each aliquot (mixture) was tested for stability of  $Cr_4S$ -to-carrier binding, by nitrogen pressure ultrafiltration through an (Amicon Corporation) YM3 (3,000-Dalton cutoff) filter, followed  
20 by washing with 5% dextrose in water. Approximately 75% retention of the (intensely blue-colored)  $Cr_4S$  ion cluster above the filter (i.e. in the retentate of more than 3,000 Dalton m.w.) was achieved by the heparin carrier. This indicated stable complexation binding of the  $Cr_4S$  ion  
25 to heparin in the presence of 834 milliosmolar dextrose (equivalent to 0.42M NaCl). Hence, the  $Cr_4S$ -heparin agent remained stably bound (complexed) under conditions equivalent to severe pathologic hyperglycemia (almost incompatible with life) and under conditions of lethal  
30 hypernatremia. Retention of  $Cr_4S$  above the filter was also observed with the dextran sulfate carrier, but was not as complete as in the presence of heparin. This partial retention in the presence of dextran sulfate carrier may be due to the close proximity of the combined  
35 molecular weight of  $Cr_4S$ -dextran sulfate carrier (ca.

10000 Daltons) to the retention cutoff value of the YM3 filter (3,000 Daltons).

Acute Toxicity testing in vivo

- 5       The Cr<sub>4</sub>S-heparin preparation (prepared as described in the preceding paragraph) was injected as an intravenous bolus into male CBA/Ln mice at ca. 3000 mg/kg, and the animals were observed for signs of acute toxicity. The animals tolerated this dose of the
- 10       preparation well immediately after injection, and were also alive, active and gaining weight normally at 1 week post-injection.

Production of Cr<sub>4</sub>S-heparin for in vivo imaging

- 15       The Cr<sub>4</sub>S ion cluster was allowed to stably complex with heparin at its stoichiometric binding ratio of ca. 30-35% (w/w) Cr<sub>4</sub>S to heparin (equivalent to about 5.4%-6.3% total chromium content (w/w to heparin). This was tested for T1 relaxivity (R1) using an IBM PC20
- 20       relaxometer (IR rf pulse sequence), and gave a 50% decrease in the water proton relaxation time at a concentration of ca. 0.33 to 1.0 mg/ml of total agent (Cr<sub>4</sub>S-heparin).

25       In Vivo MRI of tumor-bearing and control mice

- The Cr<sub>4</sub>S-heparin MRI contrast agent (from the preceding paragraph) was injected at 0.08 mmol/kg (of Cr<sub>4</sub>S ion cluster) into Balb/c mice bearing a well-differentiated, slow-growing, malignant breast tumor
- 30       induced in the mouse's lower right breast pad. Control animals consisted of uninjected normal mice and normal mice injected with Cr<sub>4</sub>S-heparin.

Specifications of MR imager, imaging conditions, and  
image processing methods from MRI tests of murine breast  
tumor, enhanced with chromium ion cluster-heparin complex  
(Cr<sub>4</sub>S-heparin)

5 Transaxial images were acquired simultaneously on three  
lightly anesthetized (i.p. pentobarbital) mice, using a  
standard clinical Diasonics whole-body, 0.35 Tesla MR  
10 imager, with the three mice oriented in the prone  
position, facing forward in the magnet and located  
centrally within a 20-cm radio frequency coil (standard  
knee coil). High resolution images were acquired over  
2.6 minute intervals, both precontrast and at several  
15 postcontrast times between 10 and 30 minutes, at an in-  
plane resolution of 0.9 x 0.9 mm and a slice thickness of  
5 mm. Five contiguous slices were acquired and the  
optimal slice photographed for maximal cross-sectional  
assessments of tumor, liver and kidney.

A pulse-sequence optimization program was run on the  
20 mice in the region of tumor, ranging from T1-weighted (TR  
= 125 msec) to T2-weighted (TR = 1800 msec) spin-echo  
sequences. The optimal spin-echo conditions were: TR of  
between 250 msec and 500 msec at an echo time (TE) of 40  
msec. Based on these results, a spin-echo pulse sequence  
25 was used of TR = 325 msec and TR = 40 msec. (Note that  
optimization of contrast enhancement under these T1-  
weighted spin-echo conditions has two important  
implications:

- 1) the Cr<sub>4</sub>S-heparin agent behaves as a "T1" contrast  
30 agent; and
- 2) the Cr<sub>4</sub>S-heparin agent is optimal for use with the more  
commonly employed, higher signal-to-noise clinical T1  
pulse sequences.

Quantitative changes in tumor image intensities in  
35 vivo, at 10 and 30 minutes after intravenous injection of  
contrast agent (Cr<sub>4</sub>S-heparin) at a dose of 0.08 mmol/kg



-91-

(based on chromium-ion complex) were assessed in the following standard fashion:

- 1) The average image intensity of central tumor region was acquired (from 64-80 image pixels each) by drawing a cursor box around the comparable central regions of tumor at precontrast, 10-minute post-contrast and 30-minute postcontrast times. (NOTE: Refer to the second, lighter grey scale, 35-mm projection slide of tumor for the exact positioning of the cursor box (this appears as a dark, irregular, rectangular-oblique line over the right-hand flank of the mouse, directly below the bright external stick-marker which was taped to the mouse skin at the site of the tumor. (See Figure 6 and Table 2.)
  - 2) The average image intensity of vertebral muscle was acquired identically (from 20-64 image pixels each). (Note: in Figure 6, note the positioning of the smaller, dark square cursor boxes located centrally and at the top (dorsum) of each mouse (in the pre, 10-min postcontrast and 30-min postcontrast panels). (See Figure 6 and Table 2.)
  - 3) Any potential artifacts in the postcontrast intensity of breast tumor which might be introduced by changes in the overall intensities of the postcontrast versus precontrast images are corrected (normalized) by the standard method (accepted experimentally and clinically), of forming the mathematical ratio of tumor-to-vertebral muscle image intensities at each imaging time (pre, 10-min post, and 30-min post). (See Table 3, which is derived from Table 2.)
- Specifications of NMR Relaxometer, relaxation conditions, and data acquisition for in vitro confirmation of tumor T1 relaxation time and pre-to-postcontrast differences in liver and kidney T1 relaxation times
- At 40 minutes after injection of contrast agent, the exact animals imaged above were sacrificed, the tumor and

-92-

organs removed and the T1 relaxation times were determined using an IBM PC20 Relaxometer (operating at 20 MHz), using a T1 inversion-recovery, 180 degrees + 90 degrees radiofrequency pulse sequence. (See Table 4.)

5       The raw data are expressed as the means of 3 to 5 individual measurements made on each organ (or tumor). the processed data are expressed as the "percent of Control" organ T1 (which is the "B" Uninjected Control animal). (Please note: A difference of about  $\pm 7\%$  is  
10       significant for each value. Hence, the decreases in kidney T1's are highly significant for both of the injected animals ("A" and "C"), but the liver T1's of these same animals are not significantly different from the uninjected control animal ("B"). Note also: Changes  
15       in the *in vitro* T1 relaxation times are inversely related to changes in the *in vivo* image intensities (enhancement of organs and tumor) at small to moderate percentage changes where T1 effects predominate and T2 effects are  
20       minor (typically at T1 percentage decreases less than about 30-40%).

#### In Vivo Tumor Imaging Results

Figure 6, which shows tumor mass in right flank, with the label of image slice "45/1" on the lower left-  
25       hand image panel, and the stick marker located just dorsal to the tumor and cursor boxes located centrally within tumor mass (at image right) and centrally within vertebral muscle (at image top). (The tumor map further clarifies orientation and dimensions.)

30       1) Precontrast: The entire tumor mass has an intermediate (grey) appearance which is relatively homogeneous. (Refer to Table 2 for the absolute values of tumor and muscle intensity.) The ratio of tumor/muscle intensity (within the cursor boxes) is  
35       1.51/1. (SEE Table 3.)

-93-

2) 10 minutes Postcontrast: The central core of the tumor mass (circumscribed by the cursor box) has increased markedly in absolute intensity, whereas the vertebral muscle has not (Table 2). The increment in central tumor intensity is such that the tumor/muscle ratio has increased to 1.98 (or by 31%). Substantial tumor architecture is now seen which was not visualized in the precontrast image (with this T1-weighted sequence - TR = 325 msec; TE = 40 msec) or any other of the pulse sequence tested - see above). The outer rim of tumor, both medial to the bottom of the cursor box, and at the lateral right border of tumor (just to the right of the bottom of the cursor box) are significantly darker than the central core. Also, a darker cystic structure with a surrounding brighter rim is present immediately below (ventral to) the bottom of the cursor box.

CORRELATION: On gross histologic examination, the three darker regions corresponded to necrotic regions of tumor, and the bright central regions corresponded to viable tumor with more extensive microvascularization.

3) 30 minutes Postcontrast: The same enhancement of tumor subregions is observed as at 10 minutes postcontrast. NOTE: The overall image intensity is slightly brighter at 30 minutes versus 10 minutes; however, as assessed by muscle intensity ratios, the muscle increment at 30 minutes is very slight (4% brighter based on absolute muscle intensity ratios of 3003/2896). Notice also that the external stick probe is visually of about the same intensity from precontrast to 30 minutes postcontrast.

NOTE: By visual inspection, the range of image pixel intensities in the central core of tumor (cursor boxes) is narrower at 10 minutes postcontrast than at 30 minutes postcontrast. This correlates with a lower standard deviation of tumor image intensity by quantitative pixel

analysis at 10 minutes postcontrast (Table 2, line 2) than at 30 minutes postcontrast (Table 2, line 3).

#### Renal and Hepatic Data

5        Figure 7 shows the renal image slices, marked 22/1, 32/1 and 44/1, respectively at the precontrast, 10-min postcontrast, and 30-min postcontrast intervals.

1) Precontrast: The faint outlines of both renal cortices appear as intermediate (grey) intensity oval  
10 outlines which are located immediately ventral and slightly lateral to the dorsal vertebral muscle, and which circumscribe the darker (black) central renal medullas.

2) 10 minutes Postcontrast: The image intensities of  
15 renal cortical regions are markedly increased in intensity, with the medullary regions being relatively darker although still absolutely increased.

3) 30 minutes Postcontrast: The image intensity of left renal cortex (the left kidney is the only one which is  
20 clearly seen in this postcontrast image) has decreased markedly relative to the 10-minute postcontrast time, but remains slightly brighter than the precontrast intensity. conversely, the left renal medulla (central region of left kidney) is quite intense, indicative of continued  
25 contrast accumulation in the renal collecting system.

Figure 8 shows a panel of 6 liver slices - 3 top and 3 bottom - at precontrast, and 10-minute postcontrast intervals.)

1) Precontrast livers are viewed in the upper row images  
30 as relatively homogeneous, intermediate-intensity (grey) masses which fill almost the entire internal abdominal volume of each mouse. The darker tube-shaped regions are larger hepatic veins and the occasional brighter regions are hepatic septae and omental fat due to caudal volume  
35 averaging (see below). The liver image of the left-hand mouse (upper row) is positioned slightly cephalad

-95-

(superior) to the optimal position for liver imaging, as indicated by the slight decrease in body width. The darker intensity of this precontrast image (relative to the other five livers) is due to slight partial volume averaging with the adjacent lung fields (which, if imaged alone, would appear black).

2) Postcontrast livers of the identical animals appear in the bottom row, as follows:

- a) left-hand animal (mouse "B") is contrast-injected (Cr,S-heparin at 0.08 mmol of Cr,S/kg) normal (nontumor) control mouse.
- b) middle animal (mouse "A") is an uninjected normal control mouse.
- c) right-hand animal (mouse "C") is a tumor mouse injected with 0.08 mmol/kg of Cr,S-heparin contrast agent.

NOTE: In an optimal 5mm-thick image slice, the liver appears relatively homogeneous and of intermediate (grey) image intensity. At both 10-minute (shown) and 30-minute (not shown) postcontrast times, dark loop densities (bowel loops) and occasional small bright nodules and ring structures (omental fat) are seen in the liver slice of the middle "A" mouse. This is not due to contrast agent (none was injected into this mouse), but is due, instead, to slight caudal mispositioning (and, hence, volume averaging) of the liver image slice. A similar but less prominent artifact is present in the left-hand "B" mouse. Minor volume averaging of this type is usually present in a multi-animal experiment because the mouse liver is only about 5mm in average height.

OVERALL RESULT: There is no significant image enhancement of liver in either of the injected mice (left "B"; or right "C") at either earlier (10-minute - shown) or later (30-minute - not shown) postcontrast times at which images were acquired.

### Preliminary Conclusions

Based on the further experimental data just described, some additional conclusions appear to be justified:

1) Enhancement of a difficult (relatively differentiated) breast carcinoma occurs acutely (10 minutes) after a low (0.08 mmol/kg) dose of the Cr<sub>3</sub>S-ion cluster-heparin complex.

2) This enhancement persists for an extended, 30-minute post-contrast interval and is not significantly decreased at 30 minutes.

NOTE: This preservation of postcontrast enhancement is markedly longer than the contrast enhancement resulting from Gd-DTPA dimeglumine (MAGNEVIST, Schering AG-Berlex) which undergoes almost complete fading (tumor clearance) by 30 minutes postcontrast.

3) Maximal image enhancement occurs in functionally viable (perfused) tumor subregions; and minimal enhancement occurs in necrotic subregions.

4) Systemic clearance of the Cr<sub>3</sub>S-heparin MRI contrast agent is predominantly by the renal route and occurs relatively quickly (first, major component requires about 30 minutes - as evidenced by the brightening and then fading of the renal cortex, with continued moderate contrast intensity in the renal medulla at 30 minutes).

5) Importantly, no major acute uptake occurs into NORMAL liver during optimal postcontrast imaging intervals.

### Interpretation and Implications.

1) Although the Cr<sub>3</sub>S-heparin contrast agent contains four Cr ions per ion complex in a superparamagnetic orientation (and, hence, exhibits intracomplex magnetic coupling), *in vivo* pulse-sequence tests indicate that this contrast agent acts as a potent T1 agent. *in vivo* dose-efficacy results indicate that this is a highly

-97-

potent agent. This result suggests that the agent has a more selective initial biodistribution compared to the existing small-molecular contrast agents (e.g., MAGNEVIST and Gd-DTPA, which exchange freely into ca. 35% of total  
5 body water and into most of the extracellular fluid, ECF).

2) Following i.v. injection, the Cr<sub>3</sub>S-heparin agent becomes sequestered rapidly in tumor interstitium but not  
10 in normal liver interstitium or parenchyma. This unique property provides for optimal body (as well as brain) imaging of tumors (including tumor within the liver) as well as potentially other body lesions (hepatitis and inflammatory/infectious lesions).

3) Tumor retention of contrast agent is prolonged  
15 relative to the rapid blood clearance (inferred from the rapid renal cortical clearance phase).

#### In Vitro T1 Measurements

*In vitro* measurements were also performed on the  
20 organs imaged in the foregoing *in vivo* imaging experiment, with results as shown in Table 4.

The T1 changes of organs freshly excised at 40 minutes postcontrast (i.v. injection) from the animals imaged above, indicate significant decreases in the T1's  
25 of kidneys for the contrast-injected "A" and "C" animals (relative to the uninjected "B" animal). However, they show no significant decreases in the T1's of livers for the injected versus uninjected animals.

These results confirm the *in vivo* imaging results  
30 and establish that the major route of clearance is renal. They also confirm that there is no significant acute clearance by normal liver.

Similarities and Differences between AgentsSimilarities of Cr,S-Heparin and Gd-DTPA-dextran

1. Increased chemical potency (increased proton T1 relaxivity) due to slower rotational correlation time of polymeric versus small molecular contrast agents
2. Restricted initial biodistribution of polymeric contrast agents *in vivo* (in ca. 10% of body water versus 35% for freely ECF-exchanging, small molecular agents)
  - a. increased *in vivo* potency
  - b. potentially decreased *in vivo* toxicity
3. Advantage of being strong T1-relaxation agents in combination with the newer, more heavily T1-weighted MRI pulse sequences
4. Improved imaging of tumors in body and brain sites, due to improved selectivity of tumor uptake
  - a. avoidance of acute uptake by normal liver (unlike standard, small contrast agents)
5. Improved detection of small tumor masses (due to increased contrast gradient at tumor margins)
6. Prolonged enhancement of tumors prior to contrast fading
  - a. patient premedication outside of imaging room
  - b. acquisition of multiple, sequential images with different pulse sequences prior to contrast fading
  - c. imaging of multiple body regions after a single dose
7. Essentially complete aqueous solubility
8. Rapid clearance by the renal route
9. Isosmotic at typical injection concentrations and doses
10. Identification and differentiation of functional tumor subregions (viable versus nonviable), due to slower interstitial diffusion of these polymeric agents relative to standard small molecular agents
  - a. improved assessment of viable tumor mass



b. noninvasive method for monitoring acute tumor-treatment effects.

Advantages of Cr,S-Heparin over Gd-DTPA-dextran

- 5 1. Cr,S-beef-lung Fraction A heparin has a lower molecular weight (less than about 10,000 Daltons) than Gd-DTPA-dextran 70 (ca. 46,000 Daltons). (This may lead to improved tumor ACCESS of Cr,S-heparin.)
2. Each Cr,S ion cluster of Cr,S-heparin contains 4
- 10 chromium ions oriented so as to produce magnetic coupling and result in a "superparamagnetic" ion complex which is ca. 1.7 times more potent than a single gadolinium ion (due to the resulting 12 unpaired electrons of Cr,S versus only 7 for gadolinium.)
- 15 3. For Cr,S, increased loading of strongly paramagnetic centers per unit length and weight of polymeric carrier (due to increased net paramagnetism in each polyatomic ion-cluster side group relative to each gadolinium-DTPA side group).
- 20 4. The heparin carrier has been established histologically (not shown) to be transported inside tumor cells, rather than just into the extracellular space surrounding tumor cells (this may contribute to prolonged contrast enhancement).
- 25 5. Possibly more rapid renal clearance of Cr,S-heparin due to smaller size of polymeric carrier.
6. Possibly increased uptake and more prolonged retention of Cr,S-heparin in tumors due to:
  - a. selective active uptake of heparin across lesional
  - 30 (tumor) endothelium; and
  - b. selective binding of heparin carrier to tissue matrix components of lesional (tumor) sites, including: fibronectin split products, laminin, collagen fragments, endogenous heparin sulfates and other matrix substances
  - 35 exposed in disease.

NOTE: Heparin which is fully complexed to  $\text{Cr}_4\text{S}$  does not produce significant in vivo anticoagulation (as assessed by the glass-induced clotting time of whole murine blood after intravenous administration of ca. 3 times the effective imaging dose).

EXAMPLE 24

10 NOVEL ACIDIC SACCHARIDE CARRIERS OF POLYATOMIC COMPLEX AND CHROMIUM  $\text{Cr}_4\text{S}$  COMPLEX -- PENTOSAN POLYSULFATE CARRIER

The polyatomic complex  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})$ , was further chemically and physically stabilized by mixing the complex and carrier together in aqueous medium with pentosan polysulfate (Sigma Chemical Co., St. Louis, MO) at a ratio of about 54 parts chromium complex to about 46 parts pentosan polysulfate (typically using stock solutions of the complex dissolved at about 1200 mg/ml and stock solutions of pentosan polysulfate dissolved at about 400 mg/ml), and the pH adjusted to about 5.0. For formulations which were further stabilized by spray drying an aqueous solution of sucrose (100 mg/ml) was added such that the final ratio of sucrose to polyatomic complex-pentosan polysulfate was about 10% by weight. For formulations which were further stabilized by lyophilization drying typically no sucrose was added. For the formulation which were dried by lyophilization further subdivisions of the formulation were made to either a) contain or b) omit the additional stabilizing (coacervating) substance, protamine, added as the sulfate salt at about 0.20% (w/w), to the preceding combined formulation components (see Table 5, "SOS" formulations 009-59A; and 009 59B'). To either of the preceding formulations, an aqueous solution of lactose was added prior to drying, such that the ratio of lactose to

polyatomic complex-pentosan polysulfate formulation was 15:1 by weight. See Table 5, "Cr<sub>4</sub>S-Carrier" for further description of physical and chemical properties.

5 a. Further stabilization by spray drying.

The preceding formulation (containing 10%, w/w of sucrose) was spray dried using a Buchi Model 190 Mini-spray drier (Brinkmann Instruments, Westbury, NY) equipped with a 7-mm atomizer nozzle, and processed at  
10 Input/Output temperatures of about 165°C/110°C, respectively. The resulting dry powder had a light, entirely blue color; and the chemical compositions of the associated Cr<sub>4</sub>S polyatomic ion cluster was retained in the native state, as assessed (following aqueous  
15 resuspension) by retention of UV peak maxima at 600nm (upper) and 438nm (lower) wavelengths, and by the retention of a UV peak-height ratio for upper/lower of 2.86 (essentially unchanged from that of the native chromium-ion cluster, equal to about 2.4 to 3.1).  
20 Submicron sizing by laser light-scattering method (Nicomp, Inc., Santa Barbara, CA), gave a nominal nanoparticulate diameter of about 112.3 nm. The Zeta potential of the particles, measured using a Delsa Model 440 surface-charge analyzer (Coulter Electronics,  
25 Hialeah, FL) was consistently negative, in the -32 to -47 millivolt (mV) range (see Table 5), indicating expression of the net negative surface charge of the pentosan polysulfate (acidic saccharide) carrier component (relative to the positively charged Cr<sub>4</sub>S ion-cluster  
30 component).

b. Further stabilization by lyophilization drying.

The preceding formulation (without added sucrose) was dried on a Virtis Model "Unitop 600SL plus  
35 Freezemobile 12" controlled shelf-temperature lyophilizer (Gardiner, NY), using an initial freezing temperature of

about -5°C, a primary-cycle temperature of about -10°C to +20°C and a secondary-cycle temperature of about +30°C. The resulting dry powder had an entirely blue color and was similar to "a." (immediately above) in all other physical and chemical parameters which were measured in common with "a." (see Table 5). Infrared spectra were obtained on formulation types either containing or omitting protamine, with the following results obtained for both types of formulation: a characteristic sharp "split-peak" absorbance peak at about 650 to 700 (cm<sup>-1</sup>) wave numbers (assigned to the intact Cr<sub>4</sub>S ion cluster -- tentatively to Cr-O and/or Cr-S bonds) and a broader peak at about 3200 (cm<sup>-1</sup>) wave numbers (as yet unassigned).

For the formulations of both "a." and "b." (above), magnetic potencies of the resulting chromium ion complex-carrier formulations were determined by water-proton relaxation analysis (R1 analysis according to: R1 = slope of line where y axis = 1/T1 versus x axis = concentration of Cr<sub>4</sub>S complex) (IBM Minispectrometer, Bruker Spectrospin CANADA Ltd. Milton, Ontario, CANADA). Comparing at equivalent concentrations of chromium-ion complex, the potencies (R1's) of the preceding formulations were equal to or greater than those of the native chromium-ion complex.

#### EXAMPLE 25

##### INITIAL TESTS OF EXCRETION ROUTE FOR FORMULATIONS OF EXAMPLE 24.

Balb/c male mice were injected intravenously with 0.1 mmol/Kg of chromium-ion cluster formulated as in Example 24 (above), the mice sacrificed at 20 minutes following injection, and their kidneys and livers analyzed for T1 relaxation-time decreases as an index of clearance of the chromium-ion cluster. Clearance occurred partially by the renal route and partially by the hepatic route. This

result suggested that a portion of the initially injected nanoparticulate material is cleared as particles (hypothetically by heparin clearance via the hepatic endothelial-hepatocyte-biliary-small bowel route) and a residual portion (probably possessing a smaller molecular radius consistent with disaggregated heparin carrier) is cleared by the glomerular filtration-renal-urinary route.

Example 26

FURTHER NOVEL ACIDIC SACCHARIDE CARRIERS OF POLYATOMIC COMPLEX AND CHROMIUM Cr<sub>3</sub>S COMPLEX -- SUCROSE OCTASULFATE CARRIER

The polyatomic complex,  $\text{Cr}_3\text{S}(\text{O}_2\text{CCH}_3)_6(\text{H}_2\text{O})_6$ , was further chemically and physically stabilized by mixing the complex and carrier together in aqueous medium with sucrose octasulfate (Toronto Research Chemicals, Inc., Downsview, Ontario, CANADA), at a ratio of about 47 parts chromium complex to about 53 parts sucrose octasulfate (typically using stock solutions of the complex dissolved at about 1200 mg/ml and the stock solutions of sucrose octasulfate dissolved at about 200 mg/ml), and the pH adjusted to about 6.6. Formulations were prepared without and with protamine (protamine sulfate) added at about 0.12% to 0.20% (w/w) to the other components of the formulation.

As in Example 24 (above), further physical stabilization was carried out by both spray drying and lyophilization drying (see Table 5).

Physical and chemical formulation parameters were measured as in Example 24 (see Table 5), with essentially similar results, except that for the parameters of submicron nanoparticulate size, which were as follows: a) the formulation without protamine had a nanoparticulate size of about 909nm (nominal mean diameter, single peak, Gaussian distribution); whereas b) the formulation with

added protamine had a significantly smaller nanoparticulate size of 11 nm (nominal mean diameter -- for its major mass of material), with two minor peaks (constituting very minor mass fractions, respectively) at 202 nm and 914 nm mean diameters (NOTE: performed by multimodal distribution analysis). By this assessment, the addition of protamine stabilized the formulation at a significantly smaller nanoparticulate size than without protamine.

10

#### EXAMPLE 27

ADDITIONAL ACIDIC SACCHARIDE CARRIERS OF POLYATOMIC COMPLEX AND CHROMIUM  $\text{Cr}_4\text{S}$  COMPLEX -- HEPARIN TOGETHER WITH LOWER (THAN PREVIOUSLY DISCLOSED) QUANTITIES OF PROTAMINE (PROTAMINE SULFATE) ADDED AS AN ADDITIONAL STABILIZER.

The polyatomic complex,  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})$ , was further chemically and physically stabilized by mixing the complex and carrier together in aqueous medium at a ratio of about 30 parts chromium complex to about 70 parts beef-lung heparin (Kabi-Pharmacia-Hepar, Inc., Franklin, OH) (typically using stock solutions of the complex dissolved at about 1200 mg/ml and stock solutions of heparin dissolved at about 400 mg/ml), with the pH adjusted to about 6.8. Protamine (protamine sulfate) was added to various of these formulation subsets, at quantities of between about 0.01% and about 0.20% (w/w) to the remaining components of the formulation. Further stabilization was carried out by lyophilization drying with lactose present at either three (3) times or fifteen (15) times the combined weight of the complex plus carrier.

#### 35 EXAMPLE 28

REPRESENTATIVE PHYSICAL AND CHEMICAL MEASUREMENTS OBTAINED FROM THE FORMULATIONS OF EXAMPLES 24 THROUGH 27 (ABOVE)

Table 5 describes the components of the formulation (chromium-ion complex--Cr<sub>3</sub>S<sub>4</sub>; Carrier -- heparin, pentosan polysulfate or sucrose octasulfate; sugar used for lyophilization drying or spray drying -- lactose and also  
5 describes the chemical and physical measurements obtained on the respective formulations: pH, chromium content (by induction-couples plasma atomic absorption analysis, ultraviolet and infrared absorption spectral analyses, particle size analysis by laser light scattering method  
10 ("Nicomp" column -- see above), and surface charge (Zeta) analysis of resulting nanoparticulate materials nominally achieved by chemical and/or physical stabilization methods (above). (Data not obtained is shown as a blank space.)

15

Figures 6a, 6b, 6c, and 6d show the following representative infrared spectra of:

20 Figure 6a: native chromium-ion cluster (with characteristic absorption maxima at 650-700 (cm<sup>-1</sup>) and 3200 (cm<sup>-1</sup>) wave numbers;

25 Figure 6b: Formulation 009-59A (see also Table 5), containing chromium-ion complex formulated with pentosan polysulfate, and lyophilized (NOTE: characteristic sharp absorbance at 650-700 (cm<sup>-1</sup>) and characteristic shoulder at 3200 (cm<sup>-1</sup>) of the native chromium-ion complex);

30 Figure 6c: Formulation 009059B (see also Table 5), containing chromium-ion complex formulated with pentosan polysulfate plus protamine (protamine sulfate) as a minor component, and lyophilized (NOTE: characteristic sharp absorbance at 650-  
35 700 (cm<sup>-1</sup>) and characteristic shoulder at 3200 (cm<sup>-1</sup>) of the native chromium-ion complex);

Figure 6d: pentosan polysulfate -- as a control for  
Figures 6a-c (above).

These infrared spectral results are representative,  
5 (with the exception of unique carrier peaks in heparin  
and sucrose octasulfate) of the IR spectra obtained from  
all other chromium ion complex-carrier formulations shown  
in Table 5 which were formulated without lactose. NOTE:  
formulations with lactose contained sufficient  
10 interference from the excess molar ratio of the lactose  
that characteristic chromium-ion complex absorbances  
could not routinely be distinguished or quantified.

TABLE 2  
15 Absolute Intensities\* of Murine Breast Tumor  
and Vertebral Muscle in MR Images (in vivo)

20	Group	Intensity of	Intensity of
		Tumor*	Muscle*
		(Mean $\pm$ 1 SD)	(Mean $\pm$ 1 SD)
	1. Precontrast	4367 $\pm$ 553	2896 $\pm$ 589
25	2. Postcontrast	5723 $\pm$ 674	2885 $\pm$ 631
	10 minutes		
	3. Postcontrast	5648 $\pm$ 801	3003 $\pm$ 436
	30 minutes		
30			

\*In arbitrary units, based on 20 to 80 image pixels



TABLE 3

Relative Intensity of Murine Breast Tumor  
in MR Images (in vivo)\*

5	<hr/>		
	Group	Intensity of	Increment
	(%)		Tumor/Muscle
10	<hr/>		
	1. Precontrast	1.51	--
	2. Postcontrast	1.98	31
	10 minutes		
15	3. Postcontrast	1.88	25
	30 minutes		
	<hr/>		

\*Data are derived from Table 2 and are based on means of  
20 to 80 image pixels

TABLE 4

T1 Relaxation Times of Freshly Excised Organs  
40 Minutes after Injection of Access MRI Contrast Agent

5

---

	<u>Animal</u>	<u>Organ</u>	<u>T1 (msec)*</u>	<u>% of Control</u>
10	A) Control, Injected	Kidney	268.5	73.2
		Liver	336.0	95.3
15	B) Control, Uninjected	Kidney	367	Control
		Liver	352.5	Control
20	C) Tumor, Injected	Kidney	269	73.3
		Liver	343	97.3
		Tumor	553	--

---

25

\*Data are means of 3 to 5 individual measurements

-109

PREP ID	TOTAL SOLIDS (mg/vial)	TABLE 3 COMPONENTS					
		(A) cR4s-% of	(B) Carrier -% of	Other -% of	Sugar -% of		
Cr4S Carrier	Spray Dried 92.5	3.36	54	2.85	PPS *	46	0.000
009-40A	Spray Dried 167.0	2.97	47	3.33	SOS *	53	0.000
009-40B	211.5	2.97	47	3.33	BLH *	53	0.0074
009-50A	Lyophilized 100.0	29.94	30	70.06	P.S.O4 **		
009-50B		7.49	30	17.51	70	0.0000	0.00
009-50C		1.87	30	4.38	70	0.0000	0.00
009-50A*		29.92	30	70.05	70	0.0125	0.01
009-50B*		7.48	30	17.51	70	0.0125	0.05
009-50C*		1.87	30	4.38	70	0.0125	0.20
009-54A	Lyophilized 100.0	29.94	30	70.06	P.S.O4 **0.00		
009-54B		29.92	30	69.99	70	0.0000	0.00
009-54C		29.88	30	69.92	70	0.1999	0.10
009-54AL		8.56	30	20.02	70	0.0000	0.20
009-54BL		8.56	30	20.01	70	0.0286	0.10
009-54CL		8.56	30	20.00	70	0.0571	0.20
009-58A1	Lyophilized 1503.0	29.94	30	69.86	P.S.O4 **		
009-58A2		7.50	30	17.49	70	0.1996	0.20
009-58B1		29.85	30	69.65	70	0.0500	0.20
009-58B2		7.49	30	17.48	70	0.4975	0.50
009-59A		54.10	54	45.90	46	0.1248	0.50
009-59B	Lyophilized 370.0	54.00	54	45.90	PPS *		
009-59A*		47.17	47	52.83	SOS *	53	0.0000
009-59B*		47.08	47	52.73	BLH *	53	0.1996
009-60A		30.00	30	70.00	P.S.O4 **		
009-60B		29.94	30	69.86	70	0.0000	0.00
009-60AL	Lyophilized 400.0	7.50	30	17.50	70	0.1996	0.20
009-60BL		7.50	30	17.49	70	0.0000	0.00

PPS = Pentosan Polysulfate; SOS = Sucrose Octasulfate; BLH = Bleef Lung Heparin  
 \*\* P.S.O4 = Protamine Sulfate; \*\*\* First number = Minspec magnet temp; second number = sample temp. > > degrees in Celsius

-110-

PREP ID	TABLE 5 (cont'd) TESTS									
	RI Value		ICP (% Theoretical)		2-Peak Max.		1-Peak Max.		L600/L440 FTIR	
	NMR (++)	pH	Input	Cr4S	Formulation	UV-VIS(nm)	UV-VIS(nm)	UV-VIS(nm)	Size (++)	Ncomp(nm)
Cr4S-Carrier	3.40	5.0	85		2.43%	438		600	2.86	+
009-40A	2.89	6.6	92.9			430		600	2.58	+
009-40B	2.80	6.6	92.9			436		602	2.92	+
009-50A	2.84	6.8	92.9		89.7	438		604	3.0	+
009-50B	3.27	6.8	92.9		87.9	438		600	3.4	+
009-50C	3.85	6.8	92.9		71.9	438		600	3.0	
009-50A'	3.14	6.8	92.9		72.4	438		602	3.0	
009-50B'	3.20	6.8	92.9		58.6	438		600	3.1	
009-50C'	3.49	6.8	92.9		58.7	438		600	2.8	
009-54A						438		602	3.02	
009-54B						438		602	2.98	
009-54C						436		602	2.83	
009-54AL						438		600	3.18	
009-54BL						438		602	3.12	
009-54CL						438		600	2.86	
009-58A1						438		604	2.88	
009-58A2						436		600	2.55	
009-58B1	2.38	6.8	99.6		106.9	438		604	2.57	+
009-58B2	3.15	6.9	99.6		76.5	434		604	2.23	+
009-59A	2.25	6.8	99.6		110.2	438		602	3.11	+
009-59B	2.30	6.8	99.6		111.0	438		602	2.74	+
009-59A'	2.50	6.8	99.6		90.2	438		602	3.02	+
009-59B'	2.44	6.8	99.6		105.2	438		602	2.89	+
009-60A		6.4	99.6		93.7	434		602	2.44	
009-60B		6.6	99.6		82.3	434		602	2.48	
009-60AL		6.5	99.6		83.4	436		600	2.86	
009-60BL		6.6	99.6		85.5	434		600	2.35	

++ Test Performed—observed characteristic peak at wave number 650; ++ All samples at pH 6.6-6.9; ++ D=distribution analysis (in descending order, small amounts in brackets); G = Gaussian distribution; ' (Ncomp) Data erratic; low count rate, large chi squared of fit error; " (Delta) Data erratic; multiple peaks, poor peak confirmation.

### Further Modifications and Variations

It will be recognized by those skilled in the art that the innovative concepts disclosed in the present application can be applied in a wide variety of contexts. 5 Moreover, the preferred implementation can be modified in a tremendous variety of ways. Accordingly, it should be understood that the modifications and variations suggested above are merely illustrative. These examples may help to show some of the scope of the inventive 10 concepts, but these examples do not nearly exhaust the full scope of variations in the disclosed novel concepts.

For example, although the presently preferred embodiment is primarily directed to imaging, the 15 selective transport advantages provided could also be used to enhance the performance of NMR spectroscopy of the human body if desired.

For another example, it is alternatively possible to 20 combine a carrier group with a small therapeutic complex. Combinations of boron (or a boroleptic group which provides a site for boron), or of cis-platinum (more precisely, cis-dichlorodiamine platinum), with a carrier group like those described above may be advantageous. 25 The active agent may be selected to provide chemotherapeutic impact, or to provide sensitization or augmentation for radiation treatment.

For one example, although the disclosed innovations 30 are particularly advantageous in selective transport to tumor sites, they can also be adapted for use with a wide variety of other types of disease or pathology, to selectively address sites where "vascular-permeability-increased" tissue exists. For example, the disclosed 35 innovations can be adapted for use in treatment or imaging (or fine-scale diagnosis) of arthritis, diabetic

angiopathy, retinitis, transplantation rejection, or other inflammatory conditions.

Similarly, the disclosed innovative ideas can also  
5 be adapted for selectively imaging sclerotic tissue, and thus may be useful in dealing with conditions such as arteriosclerosis or multiple sclerosis.

For yet another example, the disclosed innovative  
10 ideas can also be used to monitor rates of drug arrival, release, or backdiffusion.

In a further alternative, synthetic polymers other than CARBETIMER™ could be used. CARBETIMER™ is a poly-  
15 aldehyde/polyamine synthetic polymer, which provides useful transport characteristics as a carrier. Many other such synthetic polymers have been proposed, and could be used, if desired, as the polymer in the carrier.

20 It should also be noted that the carrier can be used either as a polymer or as a microsphere (or other supermolecular aggregation). Polymers are most preferably given a molecular weight in the range of less than about 10,000 Daltons, as described above; but larger  
25 polymer sizes may be advantageous for some applications. In particular, where the toxicity is very low (as with chromium), it may be advantageous to use a polymer whose molecular weight is above the renal clearance limit. In such cases, the resulting clearance time will permit the  
30 composition to be used as a "blood pool," where a low blood concentration is available over a long period of time to diffuse into a target site. (This may be particularly useful for therapeutic applications.)

35 Microspheres are even larger than the largest preferred polymer sizes. For example, a polymer of

200,000 Daltons molecular weight will have a typical maximum dimension of less than 12 nm, whereas a microsphere will have a diameter of 100 nm or more. The present invention may optionally be used with  
5 microspheres as large as 250 microns. (The larger microsphere sizes are primarily useful for embolization imaging of lung and tumor, and for imaging body cavities, such as lung, bladder, bowel, or central nervous system cavities.) Microspheres may include a surface coating  
10 which provides available reactive groups, such as hydroxyl, carbonyl, aldehyde, carboxylate, sulfate, phosphate and amine groups (singly or in combination), for binding to the complex being transported, whereas these reactive groups need not be present in the matrix  
15 of the microsphere. However, it should be noted that the polymers of the microsphere matrix should preferably be completely water-soluble, to facilitate clearance from the body.

20 It should also be noted that a composition of microspheres, with a diameter between about 0.1 micrometer and about 4.0 micrometers, and a polyatomic metal atom cluster which consists essentially of  $(Cr_4S(O_2CCH_3)_8(H_2O)_4)^{+2}$  bound to diethylenetri-  
25 aminepentaacetate-dextran, diethylenetriamine pentaacetate-hydroxyethyl starch, heparin, dextran sulfate or pentosan polysulfate, is believed to be particularly advantageous for liver imaging.

30 A further point which should be noted is that carriers (such as dextran) have been ionically coupled to an active agent, to produce drug salts; but it has not been conventional to chelate an active agent to a carrier, as is disclosed in some of the innovative  
35 examples above. As the examples above show, this further

innovative teaching is believed to provide significant advantages.

5 For superparamagnetic polyatomic structures, it should be noted that heteropolyatomic structures can be used to reduce the need for binding ligands. For example, it is known that vanadium, cobalt, or tungsten can be used as binding atoms to stabilize the relative positions of chromium atoms.

10

For another modification of the superparamagnetic polyatomic structures, it is expected that the central coordinating atom (which is sulfur in the  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$  example) could alternatively be  
15 tungsten, or even vanadium, molybdenum, cobalt, or other species.

In a further alternative, it has been found that transport of glycerol by a polymeric carrier actually  
20 increases the permeability of the vascular walls in the tumorous region. Thus, optionally, this effect can be used to further increase the selectivity of delivery of the desired agent.

25 Moreover, the varying requirements of various applications may imply a rebalancing of the various factors enumerated. For example, where it is desired to transport a metal ion which is very non-toxic, loose binding to the polymeric carrier may be perfectly  
30 acceptable. Conversely, in some cases covalent bonding may be particularly advantageous. Paired-ion embodiments may be advantageous for improved renal excretion. Polymers less than about 45K Daltons are particularly advantageous for rapid renal excretion. (Of course, in  
35 assessing the size of a polymer composition, it must be recognized that there will normally be a distribution of



sizes actually present. The references to molecular weight of polymers herein generally refer to the molecular number weight, or  $M_n$ , i.e. the peak of this distribution.)

5

The following further clarifications are included to facilitate thorough understanding and interpretation of the disclosed teachings.

- 10 1. The preferred upper limit of about twelve nanometers, for either the sizes of the carriers or the overall sizes of the carrier-metal atom agents described above, relates to experimental observations that carriers (agents) which are  
15 smaller than this approximate size are most readily and rapidly transported across endothelial (or epithelial) barriers to which a subset of the present carriers will bind as a result of their complementarity to endothelial (epithelial)  
20 determinants. Such transport (and resulting tissue access) may occur by: a) induced (active) rapid transport across physically intact (nonporous) endothelium (or epithelium); b) passive extravasation through native or induced pores  
25 (usually in sites of disease or physically compromised endothelium (or epithelium)); or c) both "a" and "b" in varying ratios, depending on the physiopathologic state of the target tissues (organs).

- 30 One of the preferred molecular weight ranges of less than about 10,000 Daltons for carriers (or carrier-metal atom agents) is based on similar considerations, and also on the experimental observations that, following intravenous administration (or other routes leading to  
35 efficient intravenous uptake and circulation):

- a) molecules larger than about 15,000 Daltons

remain predominantly within the vascular compartment except in regions of altered vascular endothelium - chemical or physical (porosity) changes - and, hence, accumulate selectively in extravascular tissue sites based on disease-induced or organ-dependent endothelial (or epithelial) binding, transport and filtration; and

- b) molecules smaller than about 45,000 Daltons are cleared efficiently by the renal route. The more restricted, preferred molecular weight range for carriers of less than about 10,000-26,000 Daltons is based on: a) the observation that pharmaceutical heparins purified from beef-lung Fraction A sources tend to fall into this general molecular weight range; and b) experiments which indicate that carriers (or complete agents) in this lower molecular weight range may undergo the very most rapid transport out of the vascular (or epithelial) compartment into underlying (potentially otherwise sequestered) tissue sites and, hence, may accumulate most efficiently in the selected tissue target sites described above. (This does not exclude, however, that the slightly larger molecular species of 26,000-45,000 Daltons - or even larger ones - could be superior for selective localization under appropriate or specialized conditions.)

In an alternative class of embodiments, lower molecular weights (as low as 1,000 Daltons) may alternatively be used. In this class of embodiments, the composition would typically be designed to bind to circulating plasma substances, and thereby reformulate itself in the body as a functionally polymeric compound

(adduct) greater than 15,000 Daltons. (Other uses of these low-MW versions may alternatively be indicated.)

2. In considering the polyatomic metal-atom complexes described above, the term "bridging ligands" extends to include both atoms and molecules and both organic and inorganic molecules. Thus, in heteropolyatomic metal-atom complexes, secondary metal atoms (paramagnetic or non-paramagnetic ones), as well as acetate, glycinate or other molecules, may serve as all or some of the bridging ligands for the primary (paramagnetic or therapeutic) metal atoms.
3. The weight ratios and molar ratios of (polyatomic) metal-atom complex-to-carrier substance are based on the following. Present examples teach that greater than about 35% (w/w) metal-atom complex-to-carrier can be achieved for Cr,S-heparin, and even higher weight ratios are referenced as possible to achieve for hydrophilic microsphere-entrapped substrates. Hence an upper limit molar ratio of about 1:2 (or about 50% w/w) is referenced here. A lower limit molar ratio of about 1:25 (or about 5% w/w) derives from: a) present examples; b) the applicant's previous patent applications (referenced above); and, importantly, c) the requirement of at least about a 5% (w/w) content of polyatomic metal-atom complex in order to achieve sufficient superparamagnetic (strongly paramagnetic) tissue proton effects for tissue visualization of intravenously injected agent, and simultaneous avoidance of unacceptably large, acute plasma volume expansion due to osmotic effects of the macromolecular carriers. (Those skilled in the art will recognize that molar ratios and weight ratios are not directly equivalent, but vary with the

molecular weights of the individual metal-atom complexes and polymeric carriers.)

4. For present purposes, the definition of "multiply  
5 paired-ion strong association" is as follows: "Two  
or more ionic charges each, of a positive and  
negative sign, located in close molecular proximity  
on the binding and bound groups, together with  
sufficient ionization of these charged groups at  
10 physiologic pH, to confer a chemical association  
which is sufficiently strong to give stable ion  
pairing in the presence of plasma or body fluids  
during localization and clearance of said metal-  
atom-carrier complexes (as elucidated in the  
15 preceding examples).
5. For the various metal atoms described above, it is  
emphasized here that numerous different metals may  
serve as either or both diagnostic or therapeutic  
20 agents (including <sup>(195)</sup>platinum, gadolinium, boron,  
gold and others). Hence, they are included together  
in the present application, and are considered all  
to be variations of a single, unified approach to  
preparing compositions of matter involving  
25 (polyatomic or other) metal-atom-complex carriers.  
In terms of various pharmaceutical applications,  
platinum is used as a chemotherapeutic as well as  
potentially a paramagnetic agent for MRI diagnosis;  
and boron and boroleptics (boron complexes) can be  
30 used either as therapeutic radiation enhancers or as  
diagnostic agents, as can gold salts. Gold salts  
and metal-atom coordinates (including, among others,  
the therapeutic antiinflammatory/antiarthritic agent  
Auraofin<sup>TM</sup>) can be administered for therapeutic  
35 purposes by formulating these salts (coordinates) as  
the metal-atom complexes described above. Hence,

the single nature and structural category of these metal-atom-carrier compositions is apparent and supported in a fashion independent of their potentially multiple diagnostic and therapeutic pharmaceutical indications.

As will be recognized by those skilled in the art, the innovative concepts described in the present application can be modified and varied over a tremendous range of applications, and accordingly their scope is not limited except by the allowed claims.

**CLAIMS:**

1. A composition of matter for enhancing induced  
internal magnetic resonance images or shifting spectra  
arising from induced internal magnetic resonance signals,  
comprising:

a polyatomic complex comprising plural atoms having  
magnetic moments which are mutually  
magnetically coupled within the polyatomic  
complex, said complex including bridging  
molecular or atomic species which are  
individually bound to at least one of said  
plural atoms having magnetic moments and other  
atoms of the polyatomic complex, said bridging  
molecular species being free of intermolecular  
bonds; and

a biocompatible, physiologically clearable carrier  
associated with the polyatomic complex, said  
carrier comprising an acidic saccharide having  
a molecular weight between about 1,000 Daltons  
and 10,000 Daltons, and binding to determinants  
of mammalian endothelia and epithelia, said  
carrier being in association with said  
polyatomic complex;

wherein said polyatomic complex and said carrier are in a  
ratio of at least about 1:20 by weight; said carrier  
being of low toxicity to mammals and containing less than  
about 5 weight percent cross-linked or microaggregated  
species.

-121-

2. The composition of claim 1 where the acidic saccharide is a sulfated saccharide.

5 3. The composition of claim 2 where the sulfated saccharide is heparin.

10 4. The composition of claim 2 where the sulfated saccharide is sucrose octasulfate, dextran sulfate, dermatan sulfate, chondroitin sulfate, or pentosan polysulfate.

15 5. The composition of claim 1 where the acidic saccharide is a carboxylated saccharide.

20 6. The composition of claim 5 where the carboxylated saccharide is hyaluronic acid.

25 7. The composition of claim 3, wherein the heparin is beef-lung heparin.

8. The composition of claim 3, wherein the heparin is Fraction A heparin.

30 9. The composition of claim 3, wherein the heparin has a mean molecular weight of about 8,000.

-122-

10. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

5 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging  
10 molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular  
15 bonds; and

a biocompatible, physiologically clearable carrier associated with the polyatomic complex, said  
20 carrier comprising Fraction A of beef-lung heparin having a molecular weight between about 6,000 Daltons and 10,000 Daltons, and binding to determinants of mammalian endothelia and  
25 epithelia; said carrier being in association with said polyatomic complex;

25 wherein said polyatomic complex and said carrier are in a ratio of at least about 1:20 by weight; said carrier being of low toxicity to mammals and containing less than  
30 about 5 weight percent cross-linked or microaggregated species.

11. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra



arising from induced internal magnetic resonance signals,  
comprising:

5 a polyatomic complex comprising plural atoms having  
magnetic moments which are mutually  
magnetically coupled within the polyatomic  
complex, said complex including bridging  
10 molecular or atomic species which are  
individually bound to at least one of said  
plural atoms having magnetic moments and other  
atoms of the polyatomic complex, said bridging  
molecular species being free of intermolecular  
bonds; and

15 a biocompatible, physiologically clearable carrier  
associated with the polyatomic complex, said  
carrier comprising heparin or a fragment  
thereof having a molecular weight between about  
1,000 Daltons and about 10,000 Daltons, and  
20 binding to determinants of mammalian endothelia  
and epithelia; said carrier substance being in  
association with said polyatomic complex;

25 wherein of said polyatomic complex and said carrier are  
in a ratio of at least about 1:20 by weight, said carrier  
containing less than about 5 weight percent cross-linked  
or microaggregated species and being of low toxicity to  
mammals.

30 12. The composition of claim 11 where the heparin is  
fraction A heparin.

13. The composition of claim 11 wherein the carrier is in association with the polyatomic complex by covalent bonding.

5

14. The composition of claim 11 wherein the heparin is native or modified.

10

15. The composition of claim 1 or 11 wherein said plural atoms include at least one of: chromium, copper, manganese, iron, platinum, cobalt, vanadium, molybdenum, tungsten, gadolinium, erbium, dysprosium, europium and holmium.

15

20

16. The composition of claim 1 or 11 wherein said polyatomic complex has the formula  $(Cr_mSR_nX_n)$ , where R is a molecular species which is at least one of formate, formaldehyde, glutaraldehyde, acetate, glycinate, succinate, acetylacetonate, malonate, propionate, glutarate, hydroxamate, oxalate, 2-bromoacetate, 2-sulfoethanoate, thiolacetate and thioglycolate; n is between 4 and 12; X is at least one of halide, sulfate, nitrate, carboxylate and phosphate stabilizing counterions; and m is between 1 and 2n.

25

30

17. The composition of claim 1 or 11 wherein the plural atoms having magnetic moments comprise Cr(III).

18. The composition of claim 1 or 11 wherein the polyatomic complex comprises four Cr(III) atoms bound to

-125-

a central tetrahedral sulfur atom and are octahedrally coordinated by bridging species.

5 19. The composition of claim 1 or 11, wherein said polyatomic complex comprises  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ .

10 20. The composition of claim 1 or 11 wherein said polyatomic complex is associated with said carrier noncovalently through a strong ionic, paired-ion, or charge interaction.

15 21. The composition of claim 20 wherein said strong interaction involves chemical coordination or chelation binding of said polyatomic complex to at least one reactive group of said carrier, each reactive group on said carrier, having a coordination number or ionic  
20 charge number between 2 and 10.

25 22. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

30 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other  
35 atoms of the polyatomic complex, said bridging

molecular species being free of intermolecular bonds; and

a biocompatible, physiologically clearable carrier associated with the polyatomic complex, said carrier comprising sucrose octasulfate and binding to determinants of mammalian endothelia and epithelia; said carrier substance being in association with said polyatomic complex;

wherein said polyatomic complex is in a ratio to said carrier of at least about 1:20 by weight, said carrier containing less than about 5 weight percent cross-linked or microaggregated species and being of low toxicity to mammals.

23. The composition of claim 22, wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ .

24. The composition of claim 22 where the sucrose octasulfate is also in association with a polyamine.

25. The composition of claim 22 wherein the sucrose octasulfate is also in association with protamine or protamine sulfate.

26. The composition of claim 25 wherein the weight ratio of protamine or protamine sulfate to polyatomic complex plus sucrose octasulfate is between about 0.12% and about 0.20%

-127-

27. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

5 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging  
10 molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular  
15 bonds; and

20 a biocompatible, physiologically clearable carrier associated with the polyatomic complex, said carrier comprising pentosan polysulfate or a fragment thereof having a molecular weight between about 1,000 Daltons and about 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia; said  
25 carrier substance being in association with said polyatomic complex;

30 wherein said polyatomic complex is in a ratio to said carrier of at least about 1:20 by weight, said carrier containing less than about 5 weight percent cross-linked or microaggregated species and being of low toxicity to mammals.

35 28. The composition of claim 27 wherein said polyatomic complex is  $\text{Cr}_3\text{S}(\text{O}_2\text{CCH}_3)_6(\text{H}_2\text{O})_4^{+2}$ .

29. The composition of claim 27 wherein the pentosan polysulfate is in association with a polyamine.

5

30. The composition of claim 27 wherein the pentosan polysulfate is in association with protamine, protamine sulfate or hexadimethrine.

10

31. The composition of claim 30 wherein the weight ratio of protamine, protamine sulfate or hexadimethrine to polyatomic complex plus pentosan polysulfate is about 0.2%.

15

32. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

20

a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular bonds; and

25

30

a biocompatible, physiologically clearable carrier associated with the polyatomic complex, said

35

-129-

carrier comprising a polyamine and said carrier binding to determinants of mammalian endothelia and epithelia; said carrier substance being in association with said polyatomic complex;

5

wherein said polyatomic complex is in a ratio to said carrier of at least about 1:20 by weight, said carrier containing less than about 5 weight percent cross-linked or microaggregated species and being of low toxicity to mammals.

10

33. The composition of claim 32 where the polyamine is hexadimethrine, protamine, protamine sulfate or a fragment thereof having a molecular weight between about 1,000 Daltons and about 10,000 Daltons,

15

34. The composition of claim 1 wherein said polyatomic complex is chemically covalently associated with said carrier.

20

35. The composition of claim 1 wherein said association is stabilized by physical or noncovalent chemical means.

25

36. The composition of claim 32, wherein said association is stabilized by heating a pre-dried polyatomic complex plus carrier combination to at least about 100 degrees Celsius in an organic solvent.

30

37. The composition of claim 36 wherein the organic solvent is acetone.

35

38. The composition of claim 36 wherein the organic solvent comprises a detergent or surfactant.

5 39. The composition of claim 36 wherein the organic solvent comprises between about 0.01% and about 25% detergent or surfactant by weight.

10 40. The composition of claim 36 wherein the organic solvent comprises a biocompatible oil.

15 41. The composition of claim 36 wherein the organic solvent comprises a polyol.

20 42. The composition of claim 41 wherein the polyol is polyethylene glycol.

43. The composition of claim 41 wherein the polyol is glycerol.

25 44. The composition of claim 1, wherein said association is stabilized by heating a pre-dried polyatomic complex and carrier combination to at least about 100 degrees Celsius in acetone, acetone comprising a polyoxyethylene sorbitan mono-oleate detergent at between about 0.01  
30 weight percent and about 25 weight percent, polyethylene glycol, glycerol or a biocompatible oil.



-131-

45. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

5 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging  
10 molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular  
15 bonds; and

a biocompatible, physiologically clearable carrier comprising protamine or protamine sulfate and heparin having a molecular weight of between  
20 about 1,000 Daltons and 10,000 Daltons and binding to determinants of mammalian endothelia and epithelia; said carrier being in association with said polyatomic complex;

25 wherein said polyatomic complex is in a ratio to said carrier of at least about 1:20 and said carrier is of low mammalian toxicity and contains less than about 5 weight percent cross-linked or microaggregated species.

30 46. The method of claim 45 wherein the heparin and protamine or protamine sulfate are in a weight ratio of between about 99.99:0.01 and 2:98.  
35

47. The composition of Claim 45, wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ .

5 48. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

10 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are  
15 individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular bonds; and

20 a biocompatible, physiologically clearable carrier comprising hexadimethrine and heparin having a molecular weight range of between about 1,000 Daltons and 10,000 Daltons and having binding  
25 to determinants of mammalian endothelia and epithelia, said carrier being bound to the polyatomic complex;

30 wherein said polyatomic complex and carrier are in a weight ratio of at least about 1:20; said carrier is of low mammalian toxicity and contains less than about 5 weight percent cross-linked or microaggregated species.

-133-

49. The composition of claim 48 wherein said hexadimethrine and heparin are in a weight ratio of between about 2:98 and 98:2.

5

50. The composition of claim 48, wherein said polyatomic complex comprises  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ .

10

51. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

15

a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are individually bound to at least one of said plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular bonds; and

20

25

a biocompatible, physiologically clearable carrier comprising starch and Fraction A of beef-lung heparin having a molecular weight range of between about 1,000 Daltons and 10,000 Daltons and binding to determinants of mammalian endothelia and epithelia; said polyatomic complex is in association with said carrier at a weight percent of at least about 5%, said carrier being in association with the polyatomic complex;

30

35

wherein said carrier is of low mammalian toxicity and contains less than about 5 weight percent cross-linked or microaggregated species.

5

52. The composition of claim 51 wherein the weight ratio of said heparin to said starch is between about 98:2 and 2:98 by weight

10

53. The composition of claim 51 wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_4^{+2}$ .

15

54. The composition of claim 1, 10, 11, 22, 27, 32, 45, 48 or 51 wherein said polyatomic complex and said carrier are further formulated to produce nanoparticles between about 1 nanometers and about 1,000 nanometers in mean diameter and said polyatomic complex is in association with a carrier comprising at least one of protamine, protamine sulfate, hexadimethrine, starch, and dextran, the acidic saccharide being at least a portion of nanoparticle surface.

20

25

55. The composition of claim 51 wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_8(\text{H}_2\text{O})_n$ , where n is 1 to 4.

30

56. The composition of claim 1, 10, 11, 22, 27, 32, 45, 48 or 51 wherein said polyatomic complex and carrier are further formulated to produce microparticles having a mean diameter between about 1 micrometer and about 15 micrometers, and said polyatomic complex is in association with a carrier comprising at least one of

35

protamine, protamine sulfate, hexadimethrine, starch, and dextran, the acidic saccharide being at least a portion of microparticle surface.

5

57. The composition of claim 56 wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_n(\text{H}_2\text{O})_n$ , where n is 1 to 4.

10

58. The composition of claim 1 wherein said polyatomic complex is  $\text{Cr}_4\text{S}(\text{O}_2\text{CCH}_3)_n(\text{H}_2\text{O})_n$ , where n is between 1 and 4, and said carrier includes at least one of a dicarboxymethyl conjugate, succinate conjugate, ethylenediaminetetraacetate conjugate or  
15 diethylenetriaminepentaacetate conjugate of said acidic saccharide.

20

59. A composition of matter comprising a metal atom complex, bound by chelation, coordination or multiply  
20 paired-ion strong association to a biocompatible, clearable carrier comprising native or modified Fraction A of beef-lung heparin having a molecular weight between about 1,000 Daltons and 10,000 Daltons, and having  
25 binding to determinants of mammalian endothelia and epithelia;

30

wherein said polyatomic complex and said carrier are in a weight ratio of at least about 1:20; and said carrier is nontoxic to mammals and contains less than about 5 weight  
30 percent cross-linked or microaggregated species.

35

60. A method for enhancing induced internal magnetic resonance images or shifting spectra arising from induced

internal magnetic resonance signals, comprising the steps of:

identifying a living vertebrate animal to be imaged;

5 introducing into the blood stream or body cavity of said animal a diagnostic imaging or enhancing agent or spectral shift agent comprising a substantially completely water-soluble  
10 polymeric carrier acidic saccharide having a molecular weight range between about 1,000 Daltons and 10,000 Daltons, and binding to determinants of mammalian endothelia and epithelia, said agent also comprising a  
15 polyatomic complex associated with said carrier, said polyatomic complex including plural atoms of magnetic moment which are mutually magnetically coupled within the polyatomic complex;

20 applying to said animal a strong magnetic field which includes a magnetic gradient;

25 applying to at least a portion of said animal an electromagnetic perturbation field at a radio frequency generally corresponding to a resonant frequency of a predetermined atomic nucleus at a magnetic field strength which falls within the range of field strengths applied to said  
30 animal by said strong magnetic field; and

measuring radio frequency emissions given off by magnetically aligned perturbed atomic nuclei, to determine a spatial map or spectral shift of

-137-

magnetic resonance characteristics within tissues of said animal.

5           61. The composition of claim 1 defined further as being stabilized by drying.

10           62. The composition of claim 61 defined further as stabilized by spray drying or lyophilization drying.

15           63. The composition of claim 1 defined further as rendered in a suitable form for administration to an animal or human by inclusion of standard pharmaceutical additives and excipients.

20           64. The composition of claim 63 wherein the additives and excipients include at least one of a sugar, salt, pH buffer, sucrose, lactose, mannitol, trehalose, monobasic and dibasic sodium or potassium phosphates, sodium hydroxide and hydrochloric acid.

25           65. The composition of claim 2 where the sulfated saccharide is pentosan polysulfate and the complex is stabilized by spray drying a preparation containing about 1 part of sucrose and about 15 parts lactose per 10 parts of carrier-polyatomic complex composition.

30

66. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra

arising from induced internal magnetic resonance signals,  
comprising:

5 a polyatomic complex comprising plural atoms having  
magnetic moments which are mutually  
magnetically coupled within the polyatomic  
complex, said complex including bridging  
10 molecular or atomic species which are  
individually bound to at least one of said  
plural atoms having magnetic moments and other  
atoms of the polyatomic complex, said bridging  
molecular species being free of intermolecular  
bonds; and

15 a biocompatible, physiologically clearable carrier  
associated with the polyatomic complex, said  
carrier comprising, in association with a  
polyamine, an acidic saccharide having a  
20 molecular weight between about 1,000 Daltons  
and 10,000 Daltons, and binding to determinants  
of mammalian endothelia and epithelia, said  
carrier being in association with said  
polyatomic complex;

25 wherein said polyatomic complex and said carrier are in a  
ratio of at least about 1:20 by weight; said carrier  
being of low toxicity to mammals and containing less than  
about 5 weight percent cross-linked or microaggregated  
species.

30  
67. The composition of claim 66 wherein the polyamine is  
protamine.



68. The composition of claim 66 wherein the polyamine is hexadimethrine.

5 69. The composition of claim 66 wherein the polyamine is protamine sulfate.

10 70. A composition of matter for enhancing induced internal magnetic resonance images or shifting spectra arising from induced internal magnetic resonance signals, comprising:

15 a polyatomic complex comprising plural atoms having magnetic moments which are mutually magnetically coupled within the polyatomic complex, said complex including bridging molecular or atomic species which are individually bound to at least one of said  
20 plural atoms having magnetic moments and other atoms of the polyatomic complex, said bridging molecular species being free of intermolecular bonds; and

25 a biocompatible, physiologically clearable carrier associated with the polyatomic complex, said carrier comprising heparin or a fragment thereof having a molecular weight between about 1,000 Daltons and about 22,000 Daltons, and  
30 binding to determinants of mammalian endothelia and epithelia; said carrier substance being in association with said polyatomic complex;

35 wherein of said polyatomic complex and said carrier are in a ratio of at least about 1:20 by weight, said carrier

containing less than about 5 weight percent cross-linked or microaggregated species and being of low toxicity to mammals.

5

71. The composition of claim 70 where the heparin is porcine heparin.

10

1/10




FIG. 1

SUBSTITUTE SHEET

2/10




FIG. 2




FIG. 3

SUBSTITUTE SHEET

3/10



BEST AVAILABLE COPY

**FIG. 4**

SUBSTITUTE SHEET

4/10




FIG. 5a




FIG. 5b

SUBSTITUTE SHEET

5/10




FIG. 5c




FIG. 5d

SUBSTITUTE SHEET

6/10




FIG. 5e

SUBSTITUTE SHEET



7/10




FIG. 6a

SUBSTITUTE SHEET

8/10




FIG. 6b

SUBSTITUTE SHEET

9/10




FIG. 6c

SUBSTITUTE SHEET

10/10




FIG. 6d